

=> file reg

FILE 'REGISTRY' ENTERED AT 10:41:26 ON 11 SEP 2003
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
 COPYRIGHT (C) 2003 American Chemical Society (ACS)

=> d his

FILE 'LREGISTRY' ENTERED AT 10:09:46 ON 11 SEP 2003

L1 STR
 L2 STR

FILE 'REGISTRY' ENTERED AT 10:20:17 ON 11 SEP 2003

L3 SCR 2043
 L4 4 S L1 AND L2 AND L3
 L5 70 S L1 AND L2 AND L3 FUL
 SAV L5 LEE223/A


FILE 'HCA' ENTERED AT 10:29:03 ON 11 SEP 2003

L6 29 S L5
 L7 38822 S ACID?(2A) (LABL? OR LABIL? OR CLEAV? OR SENS? OR DISPROP
 L8 141521 S RESIST OR RESISTS OR PHOTORESIST? OR MASK? OR PHOTOMASK
 L9 15 S L6 AND (L7 OR L8)
 L10 14 S L6 NOT L9

FILE 'REGISTRY' ENTERED AT 10:41:26 ON 11 SEP 2003

=> d l5 que stat

L1 STR



Q 9


NODE ATTRIBUTES:

NSPEC IS RC AT 9
 DEFAULT MLEVEL IS ATOM
 DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED
 NUMBER OF NODES IS 8

STEREO ATTRIBUTES: NONE
L2 STR



NODE ATTRIBUTES:

NSPEC IS RC AT 5
DEFAULT MLEVEL IS ATOM
DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:

RING(S) ARE ISOLATED OR EMBEDDED
NUMBER OF NODES IS 6

STEREO ATTRIBUTES: NONE

L3 SCR 2043
L5 70 SEA FILE=REGISTRY SSS FUL L1 AND L2 AND L3

100.0% PROCESSED 3054 ITERATIONS
SEARCH TIME: 00.00.01

70 ANSWERS


=> file hca

FILE 'HCA' ENTERED AT 10:41:48 ON 11 SEP 2003
USE IS SUBJECT TO THE TERMS OF YOUR STN/CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2003 AMERICAN CHEMICAL SOCIETY (ACS)

=> d l9 1-15 cbib abs hitstr hitind

L9 ANSWER 1 OF 15 HCA COPYRIGHT 2003 ACS on STN
139:108695 Acrylic fluoropolymers, their chemically amplified
photoresists with good vacuum UV transparency and etching
resistance, and pattern formation using them. Hatakeyama, Jun;
Harada, Yuji; Kawai, Yoshio; Sasako, Masaru; Endo, Masataka;
Kishimura, Shinji; Maeda, Kazuhiko; Otani, Michitaka; Komoritani,
Haruhiko (Shin-Etsu Chemical Industry Co., Ltd., Japan; Matsushita
Electric Industrial Co., Ltd.; Central Glass Co., Ltd.). Jpn. Kokai
Tokkyo Koho JP 2003192737 A2 20030709, 34 pp. (Japanese). CODEN:
JKXXAF. APPLICATION: JP 2001-393359 20011226.

GI



AB The invention relates to polymers having repeating units of $[CR_1(CO_2R_2)CH_2]_m$ ($R_1 = F$, C1-15-fluoroalkyl; $R_2 =$ acid-unstabilizable group; $0 < m < 1$) and I [$R_3 =$ methylene, ethylene, O, S; $R_4 = (CH_2)_aCO_2R_5$, $(CH_2)_aCR_6CO_2R_7$; $R_5, R_7 =$ acid-unstabilizable group, adhesive group, H; C1-20-alkyl, fluoroalkyl; $R_6 = H, F$, C1-20-alkyl, fluoroalkyl; $0 < n < 1$; $0 < m + n \leq 1$; $a = 0-6$]. The **photoresists** are patterned by F2 laser, Ar2 laser, or soft X ray.

IT 557771-69-4P

(chem. amplified **photoresists** with good vacuum UV transparency and etching resistance)


RN 557771-69-4 HCA

CN 2-Propenoic acid, 2-(trifluoromethyl)-, 1,1-dimethylethyl ester, polymer with .alpha.,.alpha.-bis(trifluoromethyl)-7-oxabicyclo[2.2.1]hepta-2,5-diene-2-ethanol (9CI) (CA INDEX NAME)

CM 1

CRN 557771-68-3


CMF C10 H8 F6 O2



CM 2


CRN 105935-24-8

CMF C8 H11 F3 O2



- IC ICM C08F220-22
ICS C08F232-00; C08F234-00; G03F007-039; H01L021-027
- CC 74-5 (Radiation Chemistry, Photochemistry, and Photographic and Other Reprographic Processes)
Section cross-reference(s): 38
- ST pos **photoresist** chem amplification vacuum UV; cycloolefin acrylic fluoropolymer UV laser **photoresist**; etching resistance UV **photoresist** photolithog
- IT Positive **photoresists**
(UV; chem. amplified **photoresists** with good vacuum UV transparency and etching resistance)
- IT Fluoropolymers, preparation
(acrylic; chem. amplified **photoresists** with good vacuum UV transparency and etching resistance)
- IT Photolithography
(chem. amplified **photoresists** with good vacuum UV transparency and etching resistance)
- IT 557771-65-0P 557771-66-1P 557771-67-2P 557771-69-4P
557771-71-8P
(chem. amplified **photoresists** with good vacuum UV transparency and etching resistance)
- L9 ANSWER 2 OF 15 HCA COPYRIGHT 2003 ACS on STN
139:108691 Polymers having acid-dissociable groups, chemically amplified **photoresists** with good vacuum UV transparency and etching resistance, and pattern formation using them. Hatakeyama, Jun; Harada, Yuji; Kawai, Yoshio; Sasako, Masaru; Endo, Masataka; Kishimura, Shinji; Maeda, Kazuhiko; Otani, Michitaka; Komoritani, Haruhiko (Shin-Etsu Chemical Industry Co., Ltd., Japan; Matsushita Electric Industrial Co., Ltd.; Central Glass Co., Ltd.). Jpn. Kokai Tokkyo Koho JP 2003192735 A2/20030709, 35 pp. (Japanese). CODEN: JKXXAF. APPLICATION: JP 2001-393354 20011226.

GI



- AB The invention relates to polymers having repeating units of (CR₁R₂CR₃CO₂R₄)_m (R₁, R₂ = H, F, C1-20-alkyl, fluoroalkyl; R₃ = F, C1-20-alkyl, fluoroalkyl; R₄ = acid-unstabilizable group; 0 < m ≤ 100).

$m < 1$), $(CR_1R_2CR_3OH)_n$ (R_1-3 = same as above; $0 < n < 1$), and I [R_5a , R_5b , R_6a , R_6b = H, OH, C1-20-alkyl, fluoroalkyl, $(CH_2)_dCO_2R_7$, $(CH_2)_dCR_8OR_7$; R_7 = acid-unstabilizable group, adhesive group, H, C1-20-alkyl, fluoroalkyl, etc.; R_8 = R_1 , R_2 ; $0 \leq p < 1$; $0 < m + n + p \leq 1$; $m = p \leq 0$; $c = 0, 1$; $d = 0-6$; X = methylene, ethylene, O, S]. The **photoresists** are patterned by F2 laser, Ar2 laser, or soft X ray.

IT 557112-91-1P

(chem. amplified **photoresists** with good vacuum UV transparency and etching resistance)


RN 557112-91-1 HCA

CN 2-Propenoic acid, 2-(trifluoromethyl)-, 1,1-dimethylethyl ester, polymer with .alpha.,.alpha.-bis(trifluoromethyl)-7-oxabicyclo[2.2.1]hept-5-ene-2-ethanol and 1-(trifluoromethyl)ethenyl acetate (9CI) (CA INDEX NAME)

CM 1

CRN 557104-43-5


CMF C10 H10 F6 O2



CM 2

CRN 105935-24-8


CMF C8 H11 F3 O2



CM 3


CRN 2247-91-8

CMF C5 H5 F3 O2



- IC ICM C08F220-18
ICS C08F216-02; C08F232-08; C08F234-00; G03F007-039; H01L021-027
- CC 74-5 (Radiation Chemistry, Photochemistry, and Photographic and Other Reprographic Processes)
Section cross-reference(s): 38
- ST pos **photoresist** chem amplification vacuum UV; cycloolefin acrylic fluoropolymer UV laser **photoresist**; etching resistance UV **photoresist** photolithog
- IT Positive **photoresists**
(UV; chem. amplified **photoresists** with good vacuum UV transparency and etching resistance)
- IT Fluoropolymers, preparation
(acrylic; chem. amplified **photoresists** with good vacuum UV transparency and etching-resistance)
- IT Photolithography
(chem. amplified **photoresists** with good vacuum UV transparency and etching resistance)
- IT 557112-90-0P 557112-91-1P 557112-92-2P
(chem. amplified **photoresists** with good vacuum UV transparency and etching resistance)
- L9 ANSWER 3 OF 15 HCA COPYRIGHT 2003 ACS on STN
139:108690 Chemically amplified positive **photoresists**, photolithography thereon, and polymers therefor. Harada, Yuji; Hatakeyama, Jun; Kawai, Yoshio; Sasako, Masaru; Endo, Masataka; Kishimura, Shinji; Maeda, Kazuhiko; Otani, Michitaka; Komoritani, Haruhiko (Shin-Etsu Chemical Industry Co., Ltd., Japan; Matsushita Electric Industrial Co., Ltd.; Central Glass Co., Ltd.). Jpn. Kokai Tokkyo Koho JP 2003192733 A2 20030709, 27 pp. (Japanese). CODEN: JKXXAF. APPLICATION: JP 2001-393302 20011226.

GI



- AB The **photoresists**, showing superior high sensitivity to .ltoreq.170-nm actinic rays, comprise polymers of Mw 1,000-500,000 having mer units of [CR1R2CR3(CO2R4)]a and I [R1, R2 = H, F, C1-20 (fluoro)alkyl; R3 = F, C1-20 (fluoro)alkyl; R4 = acid-labile group, coupling group, C1-20 (fluoro)alkyl; R5 = O,

S; R6a-R6d = H, OH, (CH₂)_dCR₇₂(OR₈), (CH₂)_dCO₂R₈ [R₇ = H, F, C1-20 (fluoro)alkyl; R₈ = H, **acid-labile** group, coupling group, C1-20 (fluoro)alkyl], C1-20 (fluoro)alkyl; 0 < a, b < 1; 0 < a + b .ltoreq. 1; c = 0, 1; 0 .ltoreq. d .ltoreq. 6], acid generators, and org. solvents . The **photoresists** are patternwise exposed to 100-180-nm or 1-30-nm high-energy beams (e.g., F2 laser beams, Ar2 laser beams, soft x rays) and developed (after post-exposure baking).

IT 557104-44-6P 557104-46-8P 557104-47-9P
557104-48-0P 557104-49-1P 557104-50-4P
557104-52-6P 557104-65-1P

(chem. amplified pos. **photoresists** showing superior high sensitivity to high-energy beams)


RN 557104-44-6 HCA

CN 2-Propenoic acid, 2-(trifluoromethyl)-, 1,1-dimethylethyl ester, polymer with .alpha.,.alpha.-bis(trifluoromethyl)-7-oxabicyclo[2.2.1]hept-5-ene-2-ethanol (9CI) (CA INDEX NAME)

CM 1

CRN 557104-43-5


CMF C10 H10 F6 O2



CM 2

CRN 105935-24-8

CMF C8 H11 F3 O2




RN 557104-46-8 HCA

CN 2-Propenoic acid, 2-(trifluoromethyl)-, 1,1-dimethylethyl ester, polymer with 5-[3,3,3-trifluoro-2-(methoxymethoxy)-2-(trifluoromethyl)propyl]-7-oxabicyclo[2.2.1]hept-2-ene (9CI) (CA INDEX NAME)

CM 1


COCC(C)(C)C1C=CC2C1C=CC2

CRN 105935-24-8
CMF C8 H11 F3 O2




RN	557104-47-9	HCA
CN	2-Propenoic acid, 2-(trifluoromethyl)-, 1,1-dimethylethyl ester, polymer with hexahydro-5-oxo-2,6-methanofuro[3,2-b]furan-3-yl 2-(trifluoromethyl)-2-propenoate and 5-[3,3,3-trifluoro-2- (methoxymethoxy)-2-(trifluoromethyl)propyl]-7-oxabicyclo[2.2.1]hept- 2-ene (9CI) (CA INDEX NAME)	

CRN 557104-45-7
CMF C12 H14 F6 O3




CRN 479084-31-6
CMF C11 H9 F3 O5



CM 3


CRN 105935-24-8
 CMF C8 H11 F3 O2



RN 557104-48-0 HCA
 CN 2-Propenoic acid, 2-(trifluoromethyl)-, 1,1-dimethylethyl ester,
 polymer with .alpha.,.alpha.-dimethyl-7-oxabicyclo[2.2.1]hept-5-ene-
 2-methanol (9CI) (CA INDEX NAME)


CM 1

CRN 105935-24-8
 CMF C8 H11 F3 O2



CM 2

CRN 90765-54-1
 CMF C9 H14 O2




RN 557104-49-1 HCA

CN 2-Propenoic acid, 2-(trifluoromethyl)-, 2-methyltricyclo[3.3.1.1^{3,7}]dec-2-yl ester, polymer with .alpha.,.alpha.-dimethyl-7-oxabicyclo[2.2.1]hept-5-ene-2-methanol (9CI) (CA INDEX NAME)

CM 1

CRN 188739-86-8


CMF C15 H19 F3 O2



CM 2

CRN 90765-54-1

CMF C9 H14 O2




RN 557104-50-4 HCA

CN 2-Propenoic acid, 2-(trifluoromethyl)-, 2-methyltricyclo[3.3.1.1^{3,7}]dec-2-yl ester, polymer with 5-[3,3,3-trifluoro-2-(methoxymethoxy)-2-(trifluoromethyl)propyl]-7-oxabicyclo[2.2.1]hept-2-ene (9CI) (CA INDEX NAME)

CM 1

CRN 557104-45-7


CMF C12 H14 F6 O3



CM 2

CRN 188739-86-8

CMF C15 H19 F3 O2




RN 557104-52-6 HCA

CN 2-Propenoic acid, 2-(trifluoromethyl)-, hexahydro-5-oxo-2,6-methanofuro[3,2-b]furan-3-yl ester, polymer with .alpha.,.alpha.-bis(trifluoromethyl)-7-oxabicyclo[2.2.1]hept-5-ene-2-methanol and 2-methyltricyclo[3.3.1.1^{3,7}]dec-2-yl 2-(trifluoromethyl)-2-propenoate (9CI) (CA INDEX NAME)

CM 1


CRN 557104-51-5

CMF C9 H8 F6 O2




CM 2

CRN 479084-31-6
CMF C11 H9 F3 O5



CM 3


CRN 188739-86-8
CMF C15 H19 F3 O2



RN 557104-65-1 HCA
CN 2-Propenoic acid, 2-(trifluoromethyl)-, 2-methyltricyclo[3.3.1.1^{3,7}]dec-2-yl ester, polymer with .alpha.,.alpha.-bis(trifluoromethyl)-7-oxabicyclo[2.2.1]hept-5-ene-2-ethanol (9CI) (CA INDEX NAME)

CM 1


CRN 557104-43-5
CMF C10 H10 F6 O2



CM 2


CRN 188739-86-8

CMF C15 H19 F3 O2



- IC ICM C08F220-10
ICS C08F234-00; G03F007-039; H01L021-027
- CC 74-5 (Radiation Chemistry, Photochemistry, and Photographic and Other Reprographic Processes)
Section cross-reference(s): 38
- ST fluoromethylacrylate **acid labile photoresist** polymer sensitivity; amplified
photoresist acrylic norbornene polymer oxygen incorporated
- IT Fluoropolymers, processes
(acrylic; chem. amplified pos. **photoresists** showing superior high sensitivity to high-energy beams)
- IT Photolithography
(chem. amplified pos. **photoresists** showing superior high sensitivity to high-energy beams)
- IT Positive **photoresists**
(chem. amplified; chem. amplified pos. **photoresists** showing superior high sensitivity to high-energy beams)
- IT X-ray lithography
(soft x ray; chem. amplified pos. **photoresists** showing superior high sensitivity to high-energy beams)
- IT 557104-44-6P 557104-46-8P 557104-47-9P
557104-48-0P 557104-49-1P 557104-50-4P
557104-52-6P 557104-65-1P
(chem. amplified pos. **photoresists** showing superior high sensitivity to high-energy beams)
- L9 ANSWER 4 OF 15 HCA COPYRIGHT 2003 ACS on STN
138:376396 Chemically amplified positive **photoresists** suppressing pattern shrinking for ArF excimer laser lithography. Hashimoto, Kazuhiko; Uetani, Yasunori; Fujishima, Hiroaki; Yoshida, Isao (Sumitomo Chemical Co., Ltd., Japan). Jpn. Kokai Tokkyo Koho JP 2003131381 A2 20030509, 12 pp. (Japanese). CODEN: JKXXAF. APPLICATION: JP 2001-302904 20010928. PRIORITY: JP 2001-243895 20010810.

GI



AB The **photoresists** contain alkali-insol. polymers which contain unit I [X = O, S, (m)ethylene; R₁, R₂ = H; C₁-12 alkyl, **acid-labile** group; R₃, R₄ = H, C₁-12 alkyl, **acid-labile** group, R₅CO₂R' (R₅ = direct bond, C₁-12 alkylene; R' = H, C₁-12 alkyl, **acid-labile** group), or alkyl-, lactone-, anhydride-, or ether-bearing ring; p = 0-2] and become sol. in aq. alkalis upon acid action. The polymers, which can be prepd. without metal-based catalysts, show little shrinkage upon exposure to electron beams in SEM observation.

IT 521096-27-5P 521096-28-6P 521096-29-7P
521096-30-0P

(chem. amplified pos. **photoresists** contg. alicyclic group-contg. polymers and causing no pattern shrinking in SEM observation)


RN 521096-27-5 HCA

7-Oxabicyclo[2.2.1]hept-5-ene-2-carboxylic acid, polymer with
2-methyltricyclo[3.3.1.1^{3,7}]dec-2-yl 2-methyl-2-propenoate (9CI)
(CA INDEX NAME)

CM 1

CRN 177080-67-0


CMF C15 H22 O2



CM 2

CRN 24363-23-3

CMF C7 H8 O3




RN 521096-28-6 HCA

CN 7-Oxabicyclo[2.2.1]hept-5-ene-2-carboxylic acid, polymer with
hexahydro-2-oxo-3,5-methano-2H-cyclopenta[b]furan-6-yl 2-propenoate
and 2-methyltricyclo[3.3.1.1^{3,7}]dec-2-yl 2-methyl-2-propenoate (9CI)
(CA INDEX NAME)

CM 1

CRN 242129-35-7


CMF C11 H12 O4



CM 2

CRN 177080-67-0


CMF C15 H22 O2



CM 3

CRN 24363-23-3

CMF C7 H8 O3




RN 521096-29-7 HCA

CN 7-Oxabicyclo[2.2.1]hept-5-ene-2-carboxylic acid, methyl ester,
polymer with 2-methyltricyclo[3.3.1.1^{3,7}]dec-2-yl
2-methyl-2-propenoate (9CI) (CA INDEX NAME)

CM 1

CRN 177080-67-0


CMF C15 H22 O2



CM 2

CRN 21987-33-7

CMF C8 H10 O3




RN 521096-30-0 HCA

CN 7-Oxabicyclo[2.2.1]hept-5-ene-2-carboxylic acid, polymer with
2-ethyltricyclo[3.3.1.1^{3,7}]dec-2-yl 2-methyl-2-propenoate and
hexahydro-2-oxo-3,5-methano-2H-cyclopenta[b]furan-6-yl 2-propenoate
(9CI) (CA INDEX NAME)

CM 1

CRN 242129-35-7


CMF C11 H12 O4



CM 2

CRN 209982-56-9


CMF C16 H24 O2



CM 3

CRN 24363-23-3

CMF C7 H8 O3




IC ICM G03F007-039
ICS G03F007-004; H01L021-027
CC 74-5 (Radiation Chemistry, Photochemistry, and Photographic and Other Reprographic Processes)
Section cross-reference(s): 38
ST amplified **photoresist** SEM observation pattern stability; fluoride laser transparent amplified etching **photoresist**; alicyclic acrylic polymer amplified pos **photoresist**
IT Positive **photoresists**
(chem. amplified; chem. amplified pos. **photoresists**
contg. alicyclic group-contg. polymers and causing no pattern

shrinking in SEM observation)
 IT 521096-22-0P, exo-3,6-Epoxy-1,2,3,6-tetrahydrophthalic anhydride-2-methyl-2-adamantyl 5-norbornene-2-carboxylate copolymer
 521096-24-2P 521096-26-4P 521096-27-5P
 521096-28-6P 521096-29-7P 521096-30-0P
 (chem. amplified pos. **photoresists** contg. alicyclic group-contg. polymers and causing no pattern shrinking in SEM observation)

L9 ANSWER 5 OF 15 HCA COPYRIGHT 2003 ACS on/ STN
 138:115060 Cycloalkenyl epoxy compounds, their polymers, positive **photoresists** containing them with high resolution and good adhesion to substrates, and photolithography using them. Hasegawa, Koji; Kaneo, Takeshi; Watanabe, Takeshi (Shin-Etsu Chemical Industry Co., Ltd., Japan). Jpn. Kokai Tokkyo Koho JP 2003020313 A2 20030124, 37 pp. (Japanese). CODEN: JKXXAF. APPLICATION: JP 2001-207289 20010709.

GI



AB The invention relates to epoxy compds. I (R1, R2 = H, C1-10-alkyl, etc.; R3 = C1-10-alkyl, C1-15-acyl, C1-15-alkoxycarbonyl, etc.; X = CH2, O, S; k = 0, 1; m = 0-5). The **photoresists** are sensitive to ArF excimer laser beams.

IT 488720-39-4P 488720-40-7P 488720-41-8P
 (cycloalkenyl epoxide polymers for ArF laser-sensitive high-resoln. pos. **photoresists** with good adhesion to substrates)


RN 488720-39-4 HCA

CN 2-Propenoic acid, 2-ethyltricyclo[3.3.1.1^{3,7}]dec-2-yl ester, polymer with 2,5-furandione and 5-(methoxymethyl)-7-oxabicyclo[2.2.1]hept-2-ene (9CI) (CA INDEX NAME)

CM 1

CRN 470722-58-8


CMF C8 H12 O2



CM 2

CRN 303186-14-3


CMF C15 H22 O2



CM 3

CRN 108-31-6

CMF C4 H2 O3




RN 488720-40-7 HCA

CN 2-Propenoic acid, 2-ethyltricyclo[3.3.1.13,7]dec-2-yl ester, polymer with (.alpha.,.alpha.-dimethyl-7-oxabicyclo[2.2.1]hept-5-en-2-yl)methyl acetate and 2,5-furandione (9CI) (CA INDEX NAME)

CM 1

CRN 488720-33-8


CMF C11 H16 O3



CM 2

CRN 303186-14-3


CMF C15 H22 O2



CM 3

CRN 108-31-6

CMF C4 H2 O3




RN 488720-41-8 HCA

CN 2-Propenoic acid, 2-ethyltricyclo[3.3.1.13,7]dec-2-yl ester, polymer
with 7-oxabicyclo[2.2.1]hept-5-en-2-ylmethyl acetate (9CI) (CA
INDEX NAME)

CM 1

CRN 444105-76-4


CMF C9 H12 O3



CM 2


CRN 303186-14-3

CMF C15 H22 O2



- IC ICM C08F034-00
ICS C08G061-12; G03F007-039
- CC 74-5 (Radiation Chemistry, Photochemistry, and Photographic and Other Reprographic Processes)
- ST cycloalkenyl epoxy UV excimer laser **photoresist**;
photolithog pos **resist** oxabicycloheptene polymer ArF
- IT Positive **photoresists**
(UV; cycloalkenyl epoxide polymers for ArF laser-sensitive high-resoln. pos. **photoresists** with good adhesion to substrates)
- IT Photolithography
(submicron UV; cycloalkenyl epoxide polymers for ArF laser-sensitive high-resoln. pos. **photoresists** with good adhesion to substrates)
- IT 89898-05-5P, 7-Oxabicyclo[2.2.1]hept-5-ene-2-methanol 444105-76-4P
470722-58-8P, 7-Oxabicyclo[2.2.1]hept-2-ene, 5-(methoxymethyl)-
488720-32-7P 488720-33-8P
(cycloalkenyl epoxide polymers for ArF laser-sensitive high-resoln. pos. **photoresists** with good adhesion to substrates)
- IT 488720-35-0P 488720-36-1P 488720-37-2P 488720-38-3P
488720-39-4P 488720-40-7P 488720-41-8P
488720-43-0P
(cycloalkenyl epoxide polymers for ArF laser-sensitive high-resoln. pos. **photoresists** with good adhesion to substrates)
- IT 3282-30-2, Pivaloyl chloride 21987-33-7 84752-05-6
(cycloalkenyl epoxide polymers for ArF laser-sensitive high-resoln. pos. **photoresists** with good adhesion to substrates)
- L9 ANSWER 6 OF 15 HCA COPYRIGHT 2003 ACS on STN
138:63829 **Photoresist** monomers, polymers thereof and **photoresist** compositions containing the same. Lee, Geun Su; Jung, Jae Chang; Shin, Ki Soo (S. Korea). U.S. Pat. Appl. Publ. US 2003003379 A1 20030102, 13 pp. (English). CODEN: USXXCO.
APPLICATION: US 2002-79348 20020220. PRIORITY: KR 2001-34603 20010619.

GI



AB The present invention relates to **photoresist** monomers of formula I (X1-2 = C1-10 alkylene, O,S; Y1-8 = halogen, halogen substituted alkyl; l,m = 0-3). **photoresist** polymers of it, and **photoresist** compns. contg. the same. The **photoresist** compn. has excellent etching resistance, heat resistance and adhesiveness to a wafer, and is developable in aq. tetramethylammonium hydroxide (TMAH) soln. In addn., the **photoresist** compn. has low light absorbance at 157 nm wavelength, and thus is suitable for a photolithog. process using UV light sources such as VUV (157 nm) in fabricating a minute circuit for a high integration semiconductor device.

IT 479195-52-3P

(fluoropolymer for **photoresist** compns.)


RN 479195-52-3 HCA

CN 2-Propenoic acid, 2-(trifluoromethyl)-, 1,1-dimethylethyl ester, polymer with 1,1,2,2,3,3,3a,7a-octafluoro-2,3,3a,4,7,7a-hexahydro-4,7-epoxy-1H-indene and 1-ethyl-1H-pyrrole-2,5-dione (9CI) (CA INDEX NAME)

CM 1

CRN 479195-48-7


CMF C9 H4 F8 O



CM 2


CRN 105935-24-8

CMF C8 H11 F3 O2



CM 3

CRN 128-53-0
CMF C6 H7 N O2




IC ICM G03F007-039
ICS G03F007-30; G03F007-40; G03F007-38
NCL 430018000; 430914000; 430921000; 430326000; 430330000; 430327000;
430296000; 430945000; 430942000; 430270100
CC 74-5 (Radiation Chemistry, Photochemistry, and Photographic and
Other Reprographic Processes)
Section cross-reference(s): 35, 38
ST **photoresist** monomers polymer compn photolithog UV
IT Photolithography
(UV; **photoresist** compns. contg. fluoropolymer for)
IT **Photoresists**
(**photoresist** monomers, polymers thereof and
photoresist compns. contg. same)
IT 479195-51-2P 479195-52-3P 479195-53-4P 479195-54-5P
(fluoropolymer for **photoresist** compns.)
IT 78-67-1, 2,2'-Azobisisobutyronitrile 94-36-0, Benzoylperoxide,
uses 110-05-4, tert-Butylperoxide 110-22-5, Acetylperoxide
2895-03-6, Laurylperoxide
(initiator; prepn. of polymer for **photoresist** compns.)
IT 41580-58-9 52754-92-4 57835-99-1 57840-38-7,
Triphenylsulfonium hexafluoroantimonate 57900-42-2,
Triphenylsulfonium hexafluoroarsenate 58109-40-3 62613-15-4
66003-78-9, Triphenylsulfonium triflate 81416-37-7 85342-62-7
116808-67-4, Diphenyl p-methoxyphenylsulfonium triflate
140459-13-8, Dinitrobenzyltosylate 145612-66-4 195245-87-5
255056-42-9
(photoacid generator; **photoresist** compns. contg.
fluoropolymers and)
IT 110-02-1, Thiophene 542-92-7, Cyclopentadiene, reactions
559-40-0 29797-09-9, Cyclohexadiene
(prepn. fluoropolymer for **photoresist** compns.)
IT 133205-28-4P 479195-48-7P 479195-49-8P 479195-50-1P

(prepn. fluoropolymer for **photoresist** compns.)


IT 67-68-5, Dimethylsulfoxide, uses 68-12-2, Dimethylformamide, uses 71-43-2, Benzene, uses 78-93-3, Methyl ethyl ketone, uses 108-88-3, Toluene, uses 108-94-1, Cyclohexanone, uses 109-99-9, Tetrahydrofuran, uses 120-92-3, Cyclopentanone 123-91-1, Dioxane, uses 1330-20-7, Xylene, uses (solvent; **photoresist** compns. contg.)

L9 ANSWER 7 OF 15 HCA COPYRIGHT 2003 ACS on STN
138:63824 Polymers, **resist** compositions and patterning process, novel tetrahydrofuran compounds and their preparation. Nishi, Tsunehiro; Kinsho, Takeshi; Tachibana, Seiichiro; Watanabe, Takeru; Hasegawa, Koji; Kobayashi, Tomohiro (Shin-Etsu Chemical Co., Ltd., Japan). U.S. Pat. Appl. Publ. US 2002197559 A1 20021226, 40 pp. (English). CODEN: USXXCO. APPLICATION: US 2002-126877 20020422. PRIORITY: JP 2001-124126 20010423; JP 2001-124137 20010423.

GI



I



II

AB A polymer comprises recurring units of formula I or II (R1-4 = H, alkyl; or R1,2, and R3,4 taken together may form a ring with each pair being alkylene; k = 0, 1) and having a Mw of 1,000-500,000. A **resist** compn. comprising the polymer as a base resin is sensitive to high-energy radiation, has excellent sensitivity, resolu., etching resistance, and minimized swell and lends itself to micropatterning with electron beams or deep-UV.

IT 479075-47-3P


(**photoresist** compns. and patterning process contg. novel THF polymer)

RN 479075-47-3 HCA

CN 2-Propenoic acid, 2-methyl-, 2-ethyltricyclo[3.3.1.1^{3,7}]dec-2-yl ester, polymer with 2,5-furandione and 1,3,3a,4,7,7a-hexahydro-1,1-dimethyl-4,7-epoxyisobenzofuran (9CI) (CA INDEX NAME)


CM 1

CRN 479075-38-2
CMF C10 H14 O2




CM 2

CRN 209982-56-9
CMF C16 H24 O2



CM 3


CRN 108-31-6
CMF C4 H2 O3




IC ICM G03F007-038
ICS C08G065-34; G03F007-38; G03F007-40
NCL 430270100; 528425000; 528271000; 525088000; 525165000; 430296000;
430330000; 430311000
CC 74-5 (Radiation Chemistry, Photochemistry, and Photographic and
Other Reprographic Processes)
Section cross-reference(s): 35, 38
ST **photoresist** compn patterning THF compd synthesis
IT **Photoresists**
(**photoresist** compns. and patterning process contg.
novel THF polymer)
IT 479075-39-3P 479075-41-7P 479075-42-8P 479075-44-0P
479075-45-1P 479075-46-2P **479075-47-3P** 479075-48-4P
(**photoresist** compns. and patterning process contg.
novel THF polymer)
IT 470722-61-3P 479075-38-2P 479075-40-6P

- (prepn. of novel THF compd. for **photoresist** compns. and patterning process)
- IT 98-59-9, p-Toluenesulfonyl chloride 72081-09-5 115888-24-9
479075-51-9
(prepn. of novel THF compd. for **photoresist** compns. and patterning process)
- IT 479075-49-5P 479075-50-8P
(prepn. of novel THF compd. for **photoresist** compns. and patterning process)
- L9 ANSWER 8 OF 15 HCA COPYRIGHT 2003 ACS on STN
138:63818 Novel oxanorbornene spiro derivatives and their polymers for use as **resists** for photolithographic patterning.
Hasegawa, Koji; Kaneo, Takeshi; Watanabe, Takeshi; Nishi, Tsunehiro (Shin-Etsu Chemical Industry Co., Ltd., Japan). Jpn. Kokai Tokkyo Koho JP 2002371080 A2 20021226, 39 pp. (Japanese). CODEN: JKXXAF. APPLICATION: JP 2001-179593 20010614.

GI




- AB Novel compd. I (W = CH₂, O, S; X, Y = CR₁R₂, C(O); R₁-2 = H, C₁-10 linear, branched, or cyclic alkyl with optional substitution of H with halogen; R₁ + R₂ may form aliph. ring, k may be 0) is claimed. Polymers contg. I as comonomers, **resists** mainly comprising the polymers, and photolithog. patterning of the **resists** are also claimed.
- IT 478945-98-1P
(oxanorbornene spiro deriv. (polymers) for use in chem. amplified **resists** for photolithog. patterning)
- RN 478945-98-1 HCA
CN 2-Propenoic acid, 2-ethyltricyclo[3.3.1.1^{3,7}]dec-2-yl ester, polymer with 2,5-furandione and spiro[furan-3(2H),2'-[7]oxabicyclo[2.2.1]hept[5]en]-5(4H)-one (9CI) (CA INDEX NAME)
- CM 1
- CRN 478945-85-6
CMF C9 H10 O3



CM 2

CRN 303186-14-3


CMF C15 H22 O2



CM 3

CRN 108-31-6

CMF C4 H2 O3



IC ICM C07D493-20
ICS C07D493-22; C07D495-22; C08F034-02; C08F034-04; C08G061-12;
G03F007-039; H01L021-027

CC 74-5 (Radiation Chemistry, Photochemistry, and Photographic and
Other Reprographic Processes)
Section cross-reference(s): 27, 38

ST patterning norbornene polymer chem amplified **photoresist**;
norbornene spiro compd novel; photolithog patterning norbornene
polymer chem amplified **resist**

IT **Photoresists**
(chem.-amplified; oxanorbornene spiro deriv. (polymers) for use
in chem. amplified **resists** for photolithog. patterning)

IT Polymers, preparation
(oxonorbornene spiro compds.; oxanorbornene spiro deriv.
(polymers) for use in chem. amplified **resists** for
photolithog. patterning)

IT 478945-82-3P 478945-85-6P 478945-88-9P 478945-91-4P
478945-94-7P 478945-95-8P
(oxanorbornene spiro deriv. (polymers) for use in chem. amplified
resists for photolithog. patterning)

IT 478945-83-4P 478945-86-7P 478945-89-0P 478945-92-5P
478945-96-9P **478945-98-1P** 478946-00-8P 478946-03-1P
(oxanorbornene spiro deriv. (polymers) for use in chem. amplified
resists for photolithog. patterning)
IT 109-99-9, Tetrahydrofuran, reactions 110-00-9, Furan 2170-03-8,
Itaconic acid anhydride
(oxanorbornene spiro deriv. (polymers) for use in chem. amplified
resists for photolithog. patterning)

L9 ANSWER 9 OF 15 HCA COPYRIGHT 2003 ACS on STN

137:317926 Polymer, **resist** composition and patterning process.
Nishi, Tsunehiro; Nakashima, Mutsuo; Tachibana, Seiichiro; Funatsu,
Kenji (Shin-Etsu Chemical Co., Ltd., Japan). U.S. Pat. Appl. Publ.
US 2002150835 A1 20021017, 38 pp. (English). CODEN: USXXCO.
APPLICATION: US 2002-73223 20020213. PRIORITY: JP 2001-37247
20010214; JP 2001-37262 20010214; JP 2001-37271 20010214.

AB A novel polymer is obtained by copolymerizing a (meth)acrylic acid deriv.
with a vinyl ether compound, an allyl ether compound and an
oxygen-containing alicyclic olefin compound. A **photoresist**
composition comprising the polymer as a base resin is sensitive to
high-energy radiation, has excellent sensitivity, resolution, etching
resistance, and minimized swell and lends itself to micropatterning
with electron beams or deep-UV.

IT 470722-59-9P 470722-60-2P 470722-62-4P
470722-64-6P 470722-65-7P 470722-66-8P
470722-67-9P 470722-68-0P

(polymer for **photoresist** composition and patterning process)


RN 470722-59-9 HCA

CN 2-Propenoic acid, 2-methyl-, 2-ethyltricyclo[3.3.1.1^{3,7}]dec-2-yl
ester, polymer with 5-(methoxymethyl)-7-oxabicyclo[2.2.1]hept-2-ene
(9CI) (CA INDEX NAME)

CM 1

CRN 470722-58-8

CMF C8 H12 O2




CM 2

CRN 209982-56-9

CMF C16 H24 O2

applicant




RN 470722-60-2 HCA

CN 2-Propenoic acid, 2-methyl-, 2-ethyltricyclo[3.3.1.13,7]dec-2-yl ester, polymer with 7-oxabicyclo[2.2.1]hept-5-ene-2-methanol (9CI) (CA INDEX NAME)

CM 1

CRN 209982-56-9


CMF C16 H24 O2



CM 2

CRN 89898-05-5

CMF C7 H10 O2




RN 470722-62-4 HCA

CN 2-Propenoic acid, 2-methyl-, 2-ethyltricyclo[3.3.1.13,7]dec-2-yl ester, polymer with 3'a,4',7',7'a-tetrahydrospiro[cyclopentane-1,1'(3'H)-[4,7]epoxyisobenzofuran] (9CI) (CA INDEX NAME)

CM 1


CRN 470722-61-3

CMF C12 H16 O2



CM 2


CRN 209982-56-9
 CMF C16 H24 O2



RN 470722-64-6 HCA
 CN 2-Propenoic acid, 2-methyl-, 2-ethyltricyclo[3.3.1.1.3,7]dec-2-yl ester, polymer with 1,1-dimethylethyl 7-oxabicyclo[2.2.1]hept-5-en-2-ylmethyl carbonate (9CI) (CA INDEX NAME)


CM 1

CRN 470722-63-5
 CMF C12 H18 O4



CM 2

CRN 209982-56-9
 CMF C16 H24 O2




RN 470722-65-7 HCA

CN 2-Propenoic acid, 2-methyl-, 2-ethylbicyclo[2.2.1]hept-2-yl ester,
polymer with 5-(methoxymethyl)-7-oxabicyclo[2.2.1]hept-2-ene (9CI)
(CA INDEX NAME)

CM 1

CRN 470722-58-8


CMF C8 H12 O2



CM 2

CRN 330595-98-7

CMF C13 H20 O2




RN 470722-66-8 HCA

CN 2-Propenoic acid, 2-methyl-, 1-cyclohexylcyclopentyl ester, polymer
with 5-(methoxymethyl)-7-oxabicyclo[2.2.1]hept-2-ene (9CI) (CA
INDEX NAME)

CM 1

CRN 470722-58-8


CMF C8 H12 O2



CM 2

CRN 366808-98-2

CMF C15 H24 O2




RN 470722-67-9 HCA

CN 2-Propenoic acid, 2-methyl-, polymer with 2-ethyltricyclo[3.3.1.1^{3,7}]dec-2-yl 2-methyl-2-propenoate and 5-(methoxymethyl)-7-oxabicyclo[2.2.1]hept-2-ene (9CI) (CA INDEX NAME)

CM 1

CRN 470722-58-8


CMF C8 H12 O2



CM 2

CRN 209982-56-9


CMF C16 H24 O2



CM 3

CRN 79-41-4

CMF C4 H6 O2




RN 470722-68-0 HCA

CN 2-Propenoic acid, 2-methyl-, 2-ethyltricyclo[3.3.1.1^{3,7}]dec-2-yl ester, polymer with 5-(methoxymethyl)-7-oxabicyclo[2.2.1]hept-2-ene and tetrahydro-2-oxo-3-furanyl 2-methyl-2-propenoate (9CI) (CA INDEX NAME)

CM 1

CRN 470722-58-8


CMF C8 H12 O2



CM 2

CRN 209982-56-9


CMF C16 H24 O2



CM 3

CRN 195000-66-9

CMF C8 H10 O4



IC ICM G03F007-038

ICS G03F007-20; G03F007-38; G03F007-40; G03F007-30

NCL 430270100

CC 74-5 (Radiation Chemistry, Photochemistry, and Photographic and Other Reprographic Processes)

Section cross-reference(s): 35, 38

ST **photoresist** polymer compn photolithogIT **Photoresists**(polymer for **photoresist** compn. and patterning process)

IT Photolithography

(vacuum UV; polymer for **photoresist** compn. and patterning process)

IT 470722-46-4P 470722-47-5P 470722-48-6P 470722-49-7P

470722-50-0P 470722-51-1P 470722-52-2P 470722-53-3P

470722-54-4P 470722-55-5P 470722-56-6P 470722-57-7P

470722-59-9P 470722-60-2P 470722-62-4P**470722-64-6P 470722-65-7P 470722-66-8P****470722-67-9P 470722-68-0P 470722-69-1P**

470722-70-4P 470722-71-5P 470722-72-6P 470722-73-7P

470722-74-8P 470722-75-9P 470722-76-0P

(polymer for **photoresist** compn. and patterning process)


L9 ANSWER 10 OF 15 HCA COPYRIGHT 2003 ACS on STN

137:192764 Polymer, **resist** composition and patterning process.

Nishi, Tsunehiro; Kinsho, Takeshi (Japan). U.S. Pat. Appl. Publ. US 2002115821 A1 20020822, 34 pp. (English). CODEN: USXXCO.

APPLICATION: US 2001-3117 20011206. PRIORITY: JP 2000-372406 20001207.

GI



AB The present invention relates to a polymer comprising recurring units of I and/or II ($R_{1,2}$ = H, C1-15 alkyl, acyl, alkylsulfonyl, C2-15 alkoxy carbonyl, alkoxyalkyl which may have halogen substituents; $R_{3,4}$ = H, C1-15 alkyl, alkoxy, C2-15 , alkoxyalkyl which may have halogen substituents, and $R_{3,4}$ may together bond with the carbon atom to form an aliph. ring, or $R_{3,4}$ taken together, may be an oxygen atom; $k=0$ or 1), and having a Mw of 1,000-500,000. A **resist** compn. comprising the polymer as a base resin is sensitive to high-energy radiation, has excellent sensitivity, resolu., etching resistance, and minimized swell and lends itself to micropatterning with electron beams or deep-UV.

IT 449173-04-0P 449173-05-1P

(polymer, **resist** compn. for micropatterning process)


RN 449173-04-0 HCA

CN 2-Propenoic acid, 2-ethylbicyclo[2.2.1]hept-2-yl ester, polymer with 2,5-furandione and 3a,4,7,7a-tetrahydro-2,2-dimethyl-4,7-epoxy-1,3-benzodioxole (9CI) (CA INDEX NAME)

CM 1

CRN 449173-03-9


CMF C12 H18 O2



CM 2

CRN 449172-91-2


CMF C9 H12 O3



CM 3

CRN 108-31-6

CMF C4 H2 O3




RN 449173-05-1 HCA

CN 2-Propenoic acid, 1-methyl-1-tricyclo[3.3.1.1^{3,7}]dec-1-ylethyl ester, polymer with 3a,4,7,7a-tetrahydro-4,7-epoxy-1,3-benzodioxol-2-one (9CI) (CA INDEX NAME)

CM 1

CRN 300833-10-7


CMF C16 H24 O2



CM 2

CRN 50269-96-0



CMF C7 H6 O4



IC ICM C08G065-34
 NCL 528425000
 CC 74-5 (Radiation Chemistry, Photochemistry, and Photographic and Other Reprographic Processes)
 Section cross-reference(s): 35, 38
 ST **photoresist** photolithog resin
 IT Photolithography
 (UV; polymer, **resist** compn. for micropatterning process)
 IT **Photoresists**
 (polymer, **resist** compn. for micropatterning process)
 IT 449172-89-8P 449172-90-1P 449172-92-3P 449172-94-5P
 449172-95-6P 449172-96-7P 449172-98-9P 449172-99-0P
 449173-01-7P 449173-02-8P **449173-04-0P**
449173-05-1P
 (polymer, **resist** compn. for micropatterning process)

L9 ANSWER 11 OF 15 HCA COPYRIGHT 2003 ACS on STN
 137:192763 Polymer, **resist** composition and patterning process.
 Nishi, Tsunehiro; Nakashima, Mutsuo; Kobayashi, Tomohiro (Shin-Etsu Chemical Co., Ltd., Japan). U.S. Pat. Appl. Publ. US 2002115807 A1
 20020822, 35 pp. (English). CODEN: USXXCO. APPLICATION: US
 2001-998200 20011203. PRIORITY: JP 2000-368628 20001204.

GI



AB The present invention relates to a polymer comprising recurring units of formula I or II (R1-4 = H, C1-15 alkyl, R1,2, and R3,4, taken together, may form a ring; R5,6 = H, C1-15 alkyl, acyl, alkylsulfonyl groups, C2-15 alkoxy carbonyl or alkoxyalkyl groups which may have halogen substituents; and k=0 or 1); and having a Mw of 1,000-500,000. A **resist** compn. comprising the polymer as a base resin is sensitive to high-energy radiation, has excellent sensitivity, resoln., etching resistance, and minimized swell and lends itself to micropatterning with electron beams or deep-UV.

IT **449165-82-6P**

(polymer, **resist** compn. for micropatterning process)


RN 449165-82-6 HCA

CN 2-Propenoic acid, 2-methyl-, 2-ethylbicyclo[2.2.1]hept-2-yl ester,
polymer with 2,5-furandione and 7-oxabicyclo[2.2.1]hept-5-ene-2,3-
diylbis(methylene) diacetate (9CI) (CA INDEX NAME)

CM 1

CRN 449165-64-4


CMF C12 H16 O5



CM 2

CRN 330595-98-7


CMF C13 H20 O2



CM 3

CRN 108-31-6

CMF C4 H2 O3



IC ICM C08F124-00

NCL 526266000

CC 74-5 (Radiation Chemistry, Photochemistry, and Photographic and
Other Reprographic Processes)

Section cross-reference(s): 35, 38

ST **photoresist** photolithog electron beam UV

IT Photolithography

(UV; polymer, **resist** compn. for micropatterning)

process)

IT **Photoresists**(polymer, **resist** compn. for micropatterning process)

IT 449165-65-5P 449165-69-9P 449165-73-5P 449165-76-8P



449165-78-0P 449165-80-4P **449165-82-6P** 449165-84-8P(polymer, **resist** compn. for micropatterning process)

L9 ANSWER 12 OF 15 HCA COPYRIGHT 2003 ACS on STN

137:132115 Polymer, **resist** composition and patterning process.

Nishi, Tsunehiro; Nakashima, Mutsuo; Kobayashi, Tomohiro (Shin-Etsu Chemical Co., Ltd., Japan). U.S. Pat. Appl. Publ. US 2002102493 A1 20020801, 35 pp. (English). CODEN: USXXCO. APPLICATION: US 2001-221 20011204. PRIORITY: JP 2000-368672 20001204.

GI



AB The present invention relates to a polymer comprising recurring units of I, II ($R_{1,2} = \text{H}$, C1-15 alkyl, $R_{1,2}$ taken together, may form a ring; $R_3 = \text{H}$, C1-15 alkyl, acyl or alkylsulfonyl or C2-15 alkoxy carbonyl or alkoxyalkyl which may have halogen substituents; not all R_{1-3} are hydrogen; $k = 0$ or 1) and having a Mw of 1,000-500,000.. The present invention relates to a **photoresist** compn. comprising the polymer as a base resin which is sensitive to high-energy radiation, has excellent sensitivity, resoln., etching resistance, and minimized swell and lends itself to micropatterning with electron beams or deep-UV.

IT **444105-83-3P**(polymer **photoresist** compn. for patterning process)


RN 444105-83-3 HCA

CN 2-Propenoic acid, 2-methyl-, 2-ethylbicyclo[2.2.1]hept-2-yl ester, polymer with 2,5-furandione and 7-oxabicyclo[2.2.1]hept-5-ene-2-methyl acetate (9CI) (CA INDEX NAME)

CM 1

CRN 444105-76-4


CMF C9 H12 O3



CM 2

CRN 330595-98-7


CMF C13 H20 O2



CM 3

CRN 108-31-6

CMF C4 H2 O3




IC ICM G03F007-038
 ICS G03F007-38; G03F007-40; G03F007-30
 NCL 430270100
 CC 74-5 (Radiation Chemistry, Photochemistry, and Photographic and
 Other Reprographic Processes)
 Section cross-reference(s): 35, 38
 ST **photoresists** resin photolithog
 IT Photolithography
 (UV; polymer **photoresist** compn. for patterning process)
 IT **Photoresists**
 (polymer **photoresist** compn. for patterning process)
 IT 444045-74-3P 444045-76-5P 444045-78-7P 444105-77-5P
 444105-79-7P 444105-81-1P **444105-83-3P** 444105-85-5P
 (polymer **photoresist** compn. for patterning process)

L9 ANSWER 13 OF 15 HCA COPYRIGHT 2003 ACS on STN
 136:191692 Positive-working **photoresist** compositions with high
 resolution and excellent dimensional accuracy. Mizutani, Kazuyoshi
 (Fuji Photo Film Co., Ltd., Japan). Jpn. Kokai Tokkyo Koho JP

2002055454 A2 20020220, 43 pp. (Japanese). CODEN: JKXXAF.
APPLICATION: JP 2000-244683 20000811.

GI



AB The compns. contain polymers contg. (a) structural repeating unit I (R' , R'' = H, (un)substituted alkyl, $R' + R''$ may form ring via alkylene, O, S; $R1-3$ = H, alkyl, alkoxy, halogen, trialkylsilyl, trialkylsilyloxy; n = integer of 0-3) and (b) structural repeating unit giving out acid groups on dissocn. with acid (preferable Markush given). The polymers in the compns. may also contain (c) structural repeating unit II (Z = O, $:NR_4$; R_4 = H, OH, linear or branched alkyl, O_3SR_5 ; R_5 = alkyl, trihalomethyl). Also claimed is a **photoresist** compn. contg. (A) the above stated polymers, (B) photoacid generators, and (C) org. solvents that dissolve A and B.

IT 400632-99-7 400633-03-6
(silylalkyl-contg. polymers for pos. **photoresists** showing high resolu.)


RN 400632-99-7 HCA

CN 2-Propenoic acid, 1,1-dimethylethyl ester, polymer with 2,5-furandione and 1,1,1,3,3,3-hexamethyl-2-(7-oxabicyclo[2.2.1]hept-5-en-2-yl)-2-(trimethylsilyl)trisilane (9CI) (CA INDEX NAME)

CM 1

CRN 400632-98-6


CMF C15 H34 O Si4



CM 2

CRN 1663-39-4


CMF C7 H12 O2



CM 3

CRN 108-31-6

CMF C4 H2 O3




RN 400633-03-6 HCA

CN 2-Propenoic acid, tetrahydro-2H-pyran-2-yl ester, polymer with
2,5-furandione and 1,1,1,3,3,3-hexamethyl-2-(7-oxabicyclo[2.2.1]hept-
5-en-2-yl)-2-(trimethylsilyl)trisilane (9CI) (CA INDEX NAME)

CM 1

CRN 400632-98-6


CMF C15 H34 O Si4



CM 2

CRN 52858-57-8


CMF C8 H12 O3



CM 3

CRN 108-31-6

CMF C4 H2 O3



- IC ICM G03F007-039
ICS C08F232-04; C08F234-00; C08K005-00; C08L043-04; H01L021-027
- CC 74-5 (Radiation Chemistry, Photochemistry, and Photographic and Other Reprographic Processes)
Section cross-reference(s): 38
- ST pos **photoresist** silylcyclohexane maleic anhydride copolymer
- IT Positive **photoresists**
(silylalkyl-contg. polymers for pos. **photoresists** showing high resoln.)
- IT 13891-29-7, Triphenylsulfonium tosylate 66003-76-7, Diphenyliodonium triflate 144089-15-6, Triphenylsulfonium perfluorooctanesulfonate 144317-44-2, Triphenylsulfonium nonafluorobutanesulfonate 153698-46-5, Triphenylsulfonium pentafluorophenylsulfonate 197447-16-8, Triphenylsulfonium 2,4,6-triisopropylphenylsulfonate 251463-24-8 287925-54-6, Bis(p-tert-amylphenyl)iodonium tosylate 287925-55-7, Triphenylsulfonium p-dodecylphenylsulfonate 335385-79-0
(photoacid generator; silylalkyl-contg. polymers for pos. **photoresists** showing high resoln.)
- IT 400633-31-0P 400633-35-4P
(silylalkyl-contg. polymers for pos. **photoresists** showing high resoln.)
- IT 400632-88-4 400632-90-8 400632-91-9 400632-92-0 400632-94-2
400632-99-7 400633-03-6 400633-07-0
400633-11-6 400633-15-0 400633-18-3 400633-19-4 400633-23-0
400633-26-3
(silylalkyl-contg. polymers for pos. **photoresists** showing high resoln.)
- L9 ANSWER 14 OF 15 HCA COPYRIGHT 2003 ACS on STN
- 135:280511 Positive-working **photoresist** compositions showing high resolution and high sensitivity and excellent storage stability. Sato, Kenichiro (Fuji Photo Film Co., Ltd., Japan). Jpn. Kokai Tokkyo Koho JP 2001272784 A2 20011005, 62 pp. (Japanese). CODEN: JKXXAF. APPLICATION: JP 2000-385724 20001219. PRIORITY: JP 1999-363302 19991221; JP 2000-10773 20000119; JP 2000-10774 20000119.
- AB The compns. contain (A) compds. generating acid on irradiation of actinic ray or radiation, (B) polymers contg. structural repeating unit CO2CR1R2(CR3R4)mSiR5R6R7 (R1-2 = (cyclic) alkyl; R3-4 = H, (cyclic) alkyl; R1 + R2, R3 + R4 may form cyclic alkyl; R5-7 =

(cyclic) alkyl, aryl, trialkylsilyl(oxy); m = integer of 1-6) and increasing soly. in alk. developing agents by reaction with acids, (C) org. basic compds., and (D) .gtoreq.1 of F-contg. surfactants, Si-contg. surfactants, and nonionic surfactants. Preferable structural repeating units also contained in the polymers are given in Markush. Also claimed are (1) compns. consisting of (A') acid-generating sulfonium salts $Rs1S^+ Rs2Rs3 Z^-$ ($Rs1-3 =$ (un)substituted alkyl or aryl; $Rs1 + Rs2$ may bond via single bond or bonding group; $Z^- =$ anion) and polymers B and (2) compns. consisting of acid generators A, polymers B, and certain surfactants given in the document. The compns. are useful in manuf. of semiconductor devices, printed circuits, liq. crystal panels, etc.

IT 363616-83-5P

(alk.-developing silyl-contg. polymer pos. photoresists having storage stability).


RN 363616-83-5 HCA

CN 7-Oxabicyclo[2.2.1]hept-5-ene-2-carboxylic acid, 1,1-dimethyl-3-[2,2,2-trimethyl-1,1-bis(trimethylsilyl)disilanyl]propyl ester, polymer with ethoxymethyl 2-methyl-2-propenoate and 2,5-furandione (9CI) (CA INDEX NAME)

CM 1

CRN 363616-67-5


CMF C21 H44 O3 Si4



CM 2

CRN 76392-16-0


CMF C7 H12 O3



CM 3

CRN 108-31-6

CMF C4 H2 O3




- IC ICM G03F007-039
ICS C08K005-00; C08L101-08; G03F007-004; G03F007-075; H01L021-027
- CC 74-5 (Radiation Chemistry, Photochemistry, and Photographic and Other Reprographic Processes)
Section cross-reference(s): 38
- ST pos **photoresist** alk soluble silyl contg polymer; acid generator pos **photoresist** storage stable; sulfonium salt acid generator pos **photoresist**
- IT Polysiloxanes, uses
(KP 341, surfactant; alk.-developing silyl-contg. polymer pos. **photoresists** having storage stability)
- IT Positive **photoresists**
(alk.-developing silyl-contg. polymer pos. **photoresists** having storage stability)
- IT Sulfonium compounds
(alk.-developing silyl-contg. polymer pos. **photoresists** having storage stability)
- IT Surfactants
(fluorosurfactants; alk.-developing silyl-contg. polymer pos. **photoresists** having storage stability)
- IT Surfactants
(nonionic, surfactant; alk.-developing silyl-contg. polymer pos. **photoresists** having storage stability)
- IT Fluoropolymers, uses
(surfactant; alk.-developing silyl-contg. polymer pos. **photoresists** having storage stability)
- IT 14159-45-6P 39153-56-5P 66003-76-7P 66003-78-9P 67695-82-3P
138529-81-4P 144089-15-6P 153698-46-5P 177786-98-0P
206861-54-3P 241806-75-7P 258341-95-6P 258341-99-0P
279218-73-4P 279218-74-5P 279218-75-6P 301525-08-6P
312386-77-9P 324771-13-3P 350251-56-8P 350251-57-9P
363616-18-6P
(acid generator; alk.-developing silyl-contg. polymer pos. **photoresists** having storage stability)
- IT 263713-67-3P 363616-30-2P 363616-32-4P 363616-34-6P
363616-36-8P 363616-38-0P 363616-40-4P 363616-42-6P
363616-45-9P 363616-47-1P 363616-49-3P 363616-51-7P
363616-53-9P 363616-56-2P 363616-59-5P 363616-62-0P
363616-65-3P 363616-68-6P 363616-71-1P 363616-74-4P
363616-76-6P 363616-77-7P 363616-78-8P 363616-81-3P
363616-82-4P 363616-83-5P 363616-85-7P 363616-86-8P
(alk.-developing silyl-contg. polymer pos. **photoresists** having storage stability)
- IT 484-47-9, 2,4,5-Triphenyl imidazole 1122-58-3, 4-Dimethylamino pyridine 6674-22-2, 1,8-Diazabicyclo[5.4.0]-7-undecene
(alk.-developing silyl-contg. polymer pos. **photoresists**)

- having storage stability)
- IT 96-48-0, .gamma.-Butyrolactone 96-49-1, Ethylene carbonate
97-64-3, Ethyl lactate 108-32-7, Propylene carbonate 110-43-0,
2-Heptanone 123-86-4, Butyl acetate 1320-67-8, Propylene glycol
monomethyl ether 14272-48-1, 2-Ethoxyethyl propionate
84540-57-8, Propylene glycol monomethyl ether acetate 98516-33-7,
Propylene glycol monomethyl ether propionate
(solvent; alk.-developing silyl-contg. polymer pos.
photoresists having storage stability)
- IT 9016-45-9, Polyoxyethylene nonylphenyl ether 137462-24-9, Megafac
F176 216679-67-3, Megafac R08 364039-09-8, Troysol S 336
(surfactant; alk.-developing silyl-contg. polymer pos.
photoresists having storage stability)
- L9 ANSWER-15 OF 15 HCA COPYRIGHT 2003 ACS on STN
- 134:245236 Photopolymerizable composition and chemical
amplification-type **photoresist** using it. Chung, Dong
Hang; Choi, Sang Joon; Lee, Shi Hung; Lee, Sook (Samsung Electronics
Co., Ltd., S. Korea). Jpn. Kokai Tokkyo Koho JP 2001075285 A2
20010323, 17 pp. (Japanese). CODEN: JKXXAF. APPLICATION: JP
2000-231006 20000731. PRIORITY: KR 1999-31060 19990729.
- AB The photosensitive polymer (wt. av. mol. wt. 3000-100,000) is a
copolymer of norbornene ester substituted with C1-12 aliph. alc. and
maleic acid anhydride. The polymer may have a norbornene deriv. and
an (meth)acrylic acid (ester) as other monomers. The chem.
amplification **resist** comprises the polymer and 1-15 wt.%
(based on the polymer) of a photoacid generator. The compn. shows
good adhesion with the substrate, wettability with the developer,
and good etching resistance.
- IT 329955-98-8P 329956-02-7P 329956-04-9P
329956-06-1P 329956-08-3P
(**photoresist** compn. contg. polymer from norbornene
ester and maleic anhydride and photoacid generator)
- RN 329955-98-8 HCA
- CN 7-Oxabicyclo[2.2.1]hept-5-ene-2-carboxylic acid, 3-hydroxypropyl
ester, polymer with 1,1-dimethylethyl 2-methyl-2-propenoate and
2,5-furandione (9CI) (CA INDEX NAME)

CM 1


CRN 256490-50-3

CMF C10 H14 O4




CM 2

CRN 585-07-9
 CMF C8 H14 O2



CM 3


CRN 108-31-6
 CMF C4 H2 O3



RN 329956-02-7 HCA
 CN 7-Oxabicyclo[2.2.1]hept-5-ene-2-carboxylic acid, 3-hydroxypropyl ester, polymer with 2,5-furandione and 2-methyltricyclo[3.3.1.1^{3,7}]dec-2-yl 2-propenoate (9CI) (CA INDEX NAME)


CM 1

CRN 256490-50-3
 CMF C10 H14 O4



CM 2


CRN 249562-06-9
 CMF C14 H20 O2



CM 3

CRN 108-31-6

CMF C4 H2 O3




RN 329956-04-9 HCA

CN 7-Oxabicyclo[2.2.1]hept-5-ene-2-carboxylic acid, 3-hydroxypropyl ester, polymer with 1,1-dimethylethyl bicyclo[2.2.1]hept-5-ene-2-carboxylate, 1,1-dimethylethyl 2-propenoate and 2,5-furandione (9CI)
(CA INDEX NAME)

CM 1

CRN 256490-50-3


CMF C10 H14 O4



CM 2

CRN 154970-45-3


CMF C12 H18 O2



CM 3

CRN 1663-39-4


CMF C7 H12 O2



CM 4

CRN 108-31-6

CMF C4 H2 O3




RN 329956-06-1 HCA

CN 7-Oxabicyclo[2.2.1]hept-5-ene-2-carboxylic acid, 3-hydroxypropyl ester, polymer with 1,1-dimethylethyl bicyclo[2.2.1]hept-5-ene-2-carboxylate, 2,5-furandione and rel-(1R,2R,4R)-1,7,7-trimethylbicyclo[2.2.1]hept-2-yl 2-propenoate (9CI) (CA INDEX NAME)

CM 1

CRN 256490-50-3


CMF C10 H14 O4



CM 2

CRN 154970-45-3

CMF C12 H18 O2




CM 3

CRN 5888-33-5

CMF C13 H20 O2


Relative stereochemistry.



CM 4

CRN 108-31-6

CMF C4 H2 O3




RN 329956-08-3 HCA

CN 7-Oxabicyclo[2.2.1]hept-5-ene-2-carboxylic acid, 3-hydroxypropyl ester, polymer with 1,1-dimethylethyl bicyclo[2.2.1]hept-5-ene-2-carboxylate, 2,5-furandione and 2-methyltricyclo[3.3.1.1^{3,7}]dec-2-yl 2-propenoate (9CI) (CA INDEX NAME)

CM 1

CRN 256490-50-3


CMF C10 H14 O4



CM 2

CRN 249562-06-9


CMF C14 H20 O2



CM 3

CRN 154970-45-3


CMF C12 H18 O2



CM 4

CRN 108-31-6

CMF C4 H2 O3



IC ICM G03F007-039

ICS C08F220-18; C08F222-06; C08F232-04; G03F007-004; H01L021-027

CC 74-5 (Radiation Chemistry, Photochemistry, and Photographic and

Other Reprographic Processes)

Section cross-reference(s): 38

ST **photoresist** norbornene ester maleic anhydride copolymer;
photoacid generator **photoresist**

IT **Photoresists**

(**photoresist** compn. contg. polymer from norbornene
ester and maleic anhydride and photoacid generator)

IT 102-71-6, Triethanolamine, uses 111-42-2, Diethanolamine, uses
121-44-8, Triethylamine, uses 1116-40-1, Triisobutylamine
25549-16-0, Triisooctylamine

(**photoresist** compn. contg. polymer from norbornene
ester and maleic anhydride and photoacid generator)

IT 256490-71-8P 329955-96-6P **329955-98-8P** 329956-00-5P

329956-02-7P 329956-04-9P 329956-06-1P

329956-08-3P 329956-10-7P 329956-12-9P 329956-14-1P

(**photoresist** compn. contg. polymer from norbornene
ester and maleic anhydride and photoacid generator)

IT 34684-40-7, Succinimidyl triflate 54730-01-7 66003-76-7,
Diphenyliodonium triflate 66003-78-9, Triphenylsulfonium triflate
141339-54-0 162845-55-8, Triphenylsulfonium antimonate
250345-41-6 259229-69-1

(**photoresist** compn. contg. polymer from norbornene
ester and maleic anhydride and photoacid generator)

=> d l10 1-14 cbib abs hitstr hitrn

L10 ANSWER 1 OF 14 HCA COPYRIGHT 2003 ACS on STN

136:142641 Negative working photosensitive lithographic printing plate.
Oshima, Yasuhito (Fuji Photo Film Co., Ltd., Japan). Eur. Pat.
Appl. EP 1176467 A1 20020130, 47 pp. DESIGNATED STATES: R: AT, BE,
CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT,
LV, FI, RO. (English). CODEN: EPXXDW. APPLICATION: EP 2001-116791
20010723. PRIORITY: JP 2000-228902 20000728.

AB A neg. working photosensitive lithog. printing plate is disclosed,
which comprises a support having thereon at least one photosensitive
layer contg. a polymeric binder having repeating units:
(R1MeC)-C(:O)AR2-(COOH)n (R1 = H, Me group; R2 = C3-30 hydrocarbon
group which has an alicyclic structure and a valence of n + 1; A = O,
-NR3- (R3 = H, C1-10 monovalent hydrocarbon group); and n = 1-5).
The neg. working photosensitive lithog. printing plate can attain
both high productivity and high printing durability. It is esp.
suitable for drawing with laser light.

IT **393545-89-6P 393546-28-6P**


(binder resin for neg. working photosensitive lithog. printing
plate)

RN 393545-89-6 HCA

CN 7-Oxabicyclo[2.2.1]hept-5-ene-2,3-dicarboxylic acid,
mono[2-[(1-oxo-2-propenyl)oxy]ethyl] ester, polymer with 2-propenyl
2-methyl-2-propenoate (9CI) (CA INDEX NAME)


CM 1

CRN 393545-88-5
CMF C13 H14 O7



CM 2


CRN 96-05-9
CMF C7 H10 O2



RN 393546-28-6 HCA
CN 7-Oxabicyclo[2.2.1]hept-5-ene-2,3-dicarboxylic acid,
mono[2-[(1-oxo-2-propenyl)oxy]ethyl] ester, polymer with
ethenylbenzene and 2-[(2-methyl-1-oxo-2-propenyl)oxy]ethyl
3-oxobutanoate (9CI) (CA INDEX NAME)


CM 1

CRN 393545-88-5
CMF C13 H14 O7



CM 2

CRN 21282-97-3
CMF C10 H14 O5



CM · 3

CRN 100-42-5

CMF C8 H8

 $\text{H}_2\text{C}=\text{CH}-\text{Ph}$

IT 393545-89-6P 393546-28-6P

(binder resin for neg. working photosensitive lithog. printing plate)

L10 ANSWER 2 OF 14 HCA COPYRIGHT 2003 ACS on STN

134:290081 Synthesis and antitumor and antiangiogenesis activity of polymers containing methacryloyl-2-oxy-1,2,3-propane tricarboxylic acid. Lee, Sun-Mi; Chung, Il-Doo; Lee, Neung-Ju; Ha, Chang-Sik; Lee, Chi-Ho; Cho, Won-Jei (Department of Polymer Science and Engineering, Pusan National University, Pusan, 609-735, S. Korea). Polymer International, 50(1), 119-128 (English) 2001. CODEN: PLYIEI. ISSN: 0959-8103. Publisher: John Wiley & Sons Ltd..

AB A new monomer, methacryloyl-2-oxy-1,2,3-propane tricarboxylic acid (MTCA), was prepd. from citric acid and methacrylic anhydride. Poly(methacryloyl-2-oxy-1,2,3-propane tricarboxylic acid) and poly(methacryloyl-2-oxy-1,2,3-propane tricarboxylic acid)-co-(maleic anhydride) were prepd. by radical polymn. Terpoly(methacryloyl-2-oxy-1,2,3-propane tricarboxylic acid-maleic anhydride-furan) was obtained by in situ terpolymn. of MTCA and exo-3,6-epoxy-1,2,3,6-tetrahydrophthalic anhydride. The prepd. samples were identified by FTIR, ¹H- and ¹³C-NMR spectroscopy. The no.-av. mol. wts. of the fractionated polymers detd. by GPC were in the range 14,900-16,600 and polydispersity indexes were <1.14. The in vitro IC₅₀ values of the monomer and polymers against cancer and normal cell lines were much higher than those of 5-fluorouracil (5-FU). The in vivo antitumor activities of the prepd. samples at a dosage of 0.8 mg kg⁻¹ against mice bearing the sarcoma 180 tumor cell line decreased in the order terpolymer poly(MTCA-MAH-FUR) > poly(MTCA-co-MAH) > poly(MTCA) > MTCA > 5-FU. The prepd. samples inhibited DNA replication and angiogenetic activity more than did 5-FU.

IT 334887-02-4P

(thermal- and photopolymd.; prepn. and antitumor and antiangiogenesis activity of polymers based on methacryloyloxypropane tricarboxylic acid)

RN 334887-02-4 HCA


CN 1,2,3-Propanetricarboxylic acid, 2-[(2-methyl-1-oxo-2-propenyl)oxy]-

, polymer with rel-(3aR,4S,7R,7aS)-3a,4,7,7a-tetrahydro-4,7-epoxyisobenzofuran-1,3-dione (9CI) (CA INDEX NAME)

CM 1

CRN 334886-99-6

CMF C10 H12 O8




CM 2

CRN 6118-51-0

CMF C8 H6 O4

Relative stereochemistry.



IT 334887-02-4P

(thermal- and photopolymd.; prepn. and antitumor and antiangiogenesis activity of polymers based on methacryloyloxypropane tricarboxylic acid)

L10 ANSWER 3 OF 14 HCA COPYRIGHT 2003 ACS on STN

133:282165 Novel routes to polyelectrolytes and reactive polymers via ROMP. Schitter, R. M. E.; Jocham, D.; Stelzer, F.; Moszner, N.; Volkel, Th. (Institut fuer Chemische Technologie organ. Stoffe, TU-Graz, Graz, A-8010, Austria). Journal of Applied Polymer Science, 78(1), 47-60 (English) 2000. CODEN: JAPNAB. ISSN: 0021-8995. Publisher: John Wiley & Sons, Inc..

AB Various derivs. of norbornene and 7-oxanorbornenedicarboxylic acid have been synthesized and polymd. via Ring Opening Metathesis

Polymn. (ROMP). The introduction of tetrahydropyranyl moieties as protection groups opened a way for the synthesis of polyelectrolytes through well-defined transition metal alkylidene catalysts that are usually deactivated by reactions with acidic protons. The incorporation of methacrylate groups in the polycarboxylic acids was achieved either by copolymn. of methacrylate functionalized norbornene or 7-oxanorbornene derivs., or by the polymer analogs reaction of the polycarboxylic acids with glycidyl methacrylate (GMA). These materials are sol. in water as well as in ethanol and undergo crosslinking reactions initiated by UV light.

IT 299423-40-8P 299423-41-9P 299423-42-0P
299432-78-3P

(novel routes to prep. polyelectrolytes and reactive polymers via ring-opening metathetic polymn. and structure and mol. wt. of resulting polymers)

RN 299423-40-8 HCA


CN 7-Oxabicyclo[2.2.1]hept-5-ene-2,3-dicarboxylic acid,
mono[2-[(2-methyl-1-oxo-2-propenyl)oxy]ethyl] ester,
(1R,2S,3R,4S)-rel-, homopolymer (9CI) (CA INDEX NAME)

CM 1

CRN 299423-38-4

CMF C14 H16 O7

Relative stereochemistry.



RN 299423-41-9 HCA


CN 7-Oxabicyclo[2.2.1]hept-5-ene-2,3-dicarboxylic acid, methyl
2-[(2-methyl-1-oxo-2-propenyl)oxy]ethyl ester, (1R,2S,3R,4S)-rel-,
homopolymer (9CI) (CA INDEX NAME)

CM 1

CRN 299423-39-5

CMF C15 H18 O7

Relative stereochemistry.



RN 299423-42-0 HCA


CN 7-Oxabicyclo[2.2.1]hept-5-ene-2,3-dicarboxylic acid, bis(tetrahydro-2H-pyran-2-yl) ester, (1R,2S,3R,4S)-rel-, polymer with rel-methyl (1R,2S,3R,4S)-2-[(2-methyl-1-oxo-2-propenyl)oxy]ethyl 7-oxabicyclo[2.2.1]hept-5-ene-2,3-dicarboxylate (9CI) (CA INDEX NAME)

CM 1

CRN 299423-39-5

CMF C15 H18 O7

Relative stereochemistry.




CM 2

CRN 220802-23-3

CMF C18 H24 O7


Relative stereochemistry.



RN 299432-78-3 HCA-
 CN 7-Oxabicyclo[2.2.1]hept-5-ene-2,3-dicarboxylic acid,
 (1R,2S,3R,4S)-rel-, homopolymer, 2-hydroxy-3-[(2-methyl-1-oxo-2-propenyl)oxyl]propyl ester (9CI) (CA INDEX NAME)

CM 1

CRN 5919-74-4
 CMF C7 H12 O4




CM 2

CRN 132955-58-9
 CMF (C8 H8 O5)x
 CCI PMS

CM 3

CRN 28871-62-7
 CMF C8 H8 O5

Relative stereochemistry.




IT 299423-40-8P 299423-41-9P 299423-42-0P
299432-78-3P

(novel routes to prep. polyelectrolytes and reactive polymers via ring-opening metathetic polymn. and structure and mol. wt. of resulting polymers)

L10 ANSWER 4 OF 14 HCA COPYRIGHT 2003 ACS on STN

130:272052 Dental materials based on oligomers or polymers obtained by ring-opening metathesis polymerization (ROMP). Bissinger, Peter (Espe Dental A.G., Germany). Eur. Pat. Appl. EP 904767 A2 19990331, 15 pp. DESIGNATED STATES: R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO. (German). CODEN: EPXXDW. APPLICATION: EP 1998-118366 19980929. PRIORITY: DE 1997-19742980-19970929.

GI



AB Dental filling materials, cements, inlays, veneers, etc., are prepd. from oligomers or polymers (:CHR1CH:)m or I [R1 = (substituted) C2-10 alkylene, alkenylene, or epoxyalkylene, (substituted) C6-15 arylene, di- or tetrahydrofuran-2,5-dione-3,4-diyl; R2, R3 = H, C1-15 alkyl, CO2R6, CONHR6, PO3H2,, SO3H, OH; R6 = H, (O- or N-contg.) alkyl or aryl] by ROMP with catalysis by transition metal org. compds. The starting oligomers or polymers addnl. contain groups which can be subjected to radical polymn. (leading to materials showing little shrinkage during hardening) or to hardening with a reactive filler (providing materials with improved mech. properties). Thus, 30 g exo-7-oxabicyclo[2.2.1]hept-5-ene-2,3-dicarboxylic anhydride was dissolved in 23.5 g hydroxyethyl methacrylate, stirred for 10 h, mixed with an aq. soln. of K2RuCl5.xH2O, and heated to 60.degree. to produce a viscous soln. which was dried under vacuum. The residue was combined with 10 g triethylene glycol dimethacrylate, and 10 g of the mixt. was mixed with bis(hydroxymethyl)tricyclo[5.2.1.0^{2,6}]decane diacrylate 10, camphorquinone 0.07, fumed silica 0.5, and quartz powder 79.5 g to produce a homogeneous paste. The paste was placed in a mold and hardened by irradiation with visible light. The product had a compression strength of 412 MPa, bending strength of 98 MPa, and shrinkage during polymn. of 1.6%.

IT 221881-15-8

(dental materials based on oligomers or polymers obtained by ROMP)

RN 221881-15-8 HCA

CN 2-Propenoic acid, 2-methyl-, 1,2-ethanediylbis(oxy-2,1-ethanediyl)



ester, polymer with 2-hydroxyethyl 2-methyl-2-propenoate,
(octahydro-4,7-methano-1H-indene-5,?-diyl)bis(methylene)
di-2-propenoate and rel-(3aR,4S,7R,7aS)-3a,4,7,7a-tetrahydro-4,7-
epoxyisobenzofuran-1,3-dione (9CI) (CA INDEX NAME)

CM 1

CRN 42594-17-2

CMF C18 H24 O4

CCI IDS




CM 2

CRN 6118-51-0

CMF C8 H6 O4


Relative stereochemistry.



CM 3


CRN 868-77-9

CMF C6 H10 O3



CM 4

CRN 109-16-0
CMF C14 H22 O6



IT 221881-15-8

(dental materials based on oligomers or polymers obtained by ROMP)

L10 ANSWER 5 OF 14 HCA COPYRIGHT 2003 ACS on STN

128:75744 Monomers for adhesive polymers. Part 1. Synthesis and radical polymerization of bicyclic monomers. Moszner, Norbert; Zeuner, Frank; Rheinberger, Volker (Ivoclar A.-G., Schaan, FL-9494, Liechtenstein). Polymer Bulletin (Berlin), 39(6), 669-676 (English) 1997. CODEN: POBUDR. ISSN: 0170-0839. Publisher: Springer-Verlag.

AB Bicyclic functionalized methacrylates were synthesized by a Diels-Alder reaction of furfuryl methacrylate with maleic anhydride and subsequent conversion to the corresponding monomers contg. carboxylic groups. The structure of the bicyclic functionalized methacrylates was confirmed by elemental anal., IR, and ¹H and ¹³C NMR. The radical photopolymer. of bicyclic monomethacrylates in DMF with 2,2'-azobisisobutyronitrile (AIBN) resulted in sol. polymers, while a bicyclic dimethacrylate resulted in a crosslinked polymer.

IT 31343-32-5P 200507-11-5P 200507-12-6P

200507-13-7P


(prepn. and radical polymer. of carboxyl-contg. bicyclic methacrylates)

RN 31343-32-5 HCA

CN 2-Propenoic acid, 2-methyl-, (3,3a,7,7a-tetrahydro-1,3-dioxo-4,7-epoxyisobenzofuran-4(1H)-yl)methyl ester, homopolymer (9CI) (CA INDEX NAME)

CM 1

CRN 7726-38-7
CMF C13 H12 O6




RN 200507-11-5 HCA

CN 7-Oxabicyclo[2.2.1]hept-5-ene-2,3-dicarboxylic acid,
1-[[2-methyl-1-oxo-2-propenyl]oxy]methyl-, homopolymer (9CI) (CA
INDEX NAME)

CM 1

CRN 200507-05-7

CMF C13 H14 O7




RN 200507-12-6 HCA

CN 7-Oxabicyclo[2.2.1]hept-5-ene-2,3-dicarboxylic acid,
1-[[2-methyl-1-oxo-2-propenyl]oxy]methyl-, 2-(2-hydroxyethyl)
ester, homopolymer (9CI) (CA INDEX NAME)

CM 1

CRN 200507-06-8

CMF C15 H18 O8




RN 200507-13-7 HCA

CN 7-Oxabicyclo[2.2.1]hept-5-ene-2,3-dicarboxylic acid,
1-[[[(2-methyl-1-oxo-2-propenyl)oxy]methyl]-, 2,2'-(1,2-ethanediyl)
ester, homopolymer (9CI) (CA INDEX NAME)

CM 1

CRN 200507-10-4

CMF C28 H30 O14



IT 31343-32-5P 200507-11-5P 200507-12-6P
200507-13-7P

(prepn. and radical polymn. of carboxyl-contg. bicyclic
methacrylates)

L10 ANSWER 6 OF 14 HCA COPYRIGHT 2003 ACS on STN

128:43560 Syntheses and biological activities of 5'-O-methacryloyl-3'-
azido-3'-deoxythymidine and its polymers. Choi, Won Moon; Lee,
Neung Ju; Lee, Young Woo; Ha, Chang Sik; Cho, Won Jei (Dep. Polymer
Sci. Eng., Pusan National Univ., Pusan, 609735, S. Korea). Polymer
Bulletin (Berlin), 39(5), 535-542 (English) 1997. CODEN: POBUDR.
ISSN: 0170-0839. Publisher: Springer-Verlag.

AB The new monomer, 5'-O-methacryloyl-3'-azido-3'-deoxythymidine
(MAZT), was synthesized from methacryloyl chloride (MAC) and
3'-azido-3'-deoxythymidine (AZT). The homopolymer of MAZT and
copolymers of MAZT with acrylic acid (AA) or exo-3,6-epoxy-1,2,3,6-

tetrahydrophthalic glycinylimide (ETGI) were synthesized by radical polymns. The structures of MAZT and polymers were confirmed identified by FT-IR and ¹H-NMR spectroscopies. The no. av. mol. wts. (.hivin.Mn) and polydispersity indexes of the synthesized polymers were in the range of 4,400-20,400 and 1.2-2.0. The in vitro cytotoxicities of polymers against K562 human leukemia and normal cell lines were greater than that of control.

IT 199998-18-0P

(prepn. and biol. activities of O-methacryloyl azidodeoxythymidine and its polymers)

RN 199998-18-0 HCA


CN Thymidine, 3'-azido-3'-deoxy-, 5'-(2-methyl-2-propenoate), polymer with (3a.alpha.,4.beta.,7.beta.,7a.alpha.)-1,3,3a,4,7,7a-hexahydro-1,3-dioxo-4,7-epoxy-2H-isoindole-2-acetic acid (9CI) (CA INDEX NAME)

CM 1

CRN 193471-62-4

CMF C14 H17 N5 O5

Absolute stereochemistry.




CM 2

CRN 182505-67-5

CMF C10 H9 N O5

Relative stereochemistry.



IT 199998-18-0P

(prepn. and biol. activities of O-methacryloyl
azidodeoxythymidine and its polymers)

L10 ANSWER 7 OF 14 HCA COPYRIGHT 2003 ACS on STN

127:253230 Functionalized bicyclic (meth)acrylates for dental adhesives.
Moszner, Norbert; Rheinberger, Völker; Vogel, Karin; Zeuner, Frank
(Ivoclar Ag, Liechtenstein). Ger. Offen. DE 19608316 A1 19970828,
18 pp. (German). CODEN: GWXXBX. APPLICATION: DE 1996-19608316
19960222.

AB Functionalized bicyclic (meth)acrylates with norbornenyl or
norbornadienyl groups are described which undergo radical polymn. at
room temp. and are excellent adhesion promoters for use in dental
adhesive compns. Thus, a dental adhesive primer contg.
exo-1-[(methacryloyloxy)methyl]-7-oxabicyclo[2.2.1]hept-5-ene-2,3-
dicarboxylic acid mono(2-hydroxyethyl) ester (I) 20, HEMA 20,
camphorquinone 0.3, diphenyliodonium hexafluorophosphate
(accelerator) 1, H₂O 28.7, and EtOH 30 wt.% was applied to a dentin
surface for 30 s and blown off, followed by a photohardening bonding
compn. (bis-GMA 60, triethylene glycol dimethacrylate 39.26,
cyanoethyl(methyl)aniline 0.5, and camphorquinone 0.24 wt.%) and
illumination, and finally a photohardening filling compn. (Tetric).
I was prepd. by Diels-Alder addn. of maleic anhydride to furfuryl
methacrylate, followed by hydrolysis of the anhydride and
esterification with ethylene glycol.

IT 195376-93-3P 195376-94-4P 195442-95-6P
195540-15-9P

(functionalized bicyclic (meth)acrylates for dental adhesives)

RN 195376-93-3 HCA


CN 2-Propenoic acid, 2-methyl-, [(3aR,4S,7R,7aS)-3,3a,7,7a-tetrahydro-
1,3-dioxo-4,7-epoxyisobenzofuran-4(1H)-yl]methyl ester, homopolymer
(9CI) (CA INDEX NAME)

CM 1

CRN 195376-88-6

CMF C13 H12 O6

Absolute stereochemistry.




RN 195376-94-4 HCA
 CN 7-Oxabicyclo[2.2.1]hept-5-ene-2,3-dicarboxylic acid,
 1-[[[(2-methyl-1-oxo-2-propenyl)oxy]methyl]-, [1S-(exo,exo)]-,
 homopolymer (9CI) (CA INDEX NAME)

CM 1

CRN 195376-89-7

CMF C13 H14 O7

Absolute stereochemistry.




RN 195442-95-6 HCA
 CN 7-Oxabicyclo[2.2.1]hept-5-ene-2,3-dicarboxylic acid,
 1-[[[(2-methyl-1-oxo-2-propenyl)oxy]methyl]-, mono(2-hydroxyethyl)
 ester, [1S-(exo,exo)]-, polymer with 2-hydroxyethyl
 2-methyl-2-propenoate (9CI) (CA INDEX NAME)

CM 1

CRN 868-77-9

CMF C6 H10 O3



CM 2

CRN 195442-94-5

CMF C15 H18 O8


CCI IDS

CM 3

CRN 195376-89-7

CMF C13 H14 O7


Absolute stereochemistry.



CM 4

CRN 107-21-1

CMF C2 H6 O2



RN 195540-15-9 HCA

CN 7-Oxabicyclo[2.2.1]hept-5-ene-2,3-dicarboxylic acid,
1-[[[(2-methyl-1-oxo-2-propenyl)oxy]methyl]-, mono(2-hydroxyethyl)
ester, [1S-(exo,exo)]-, homopolymer (9CI) (CA INDEX NAME)

CM 1


CRN 195442-94-5

CMF C15 H18 O8
CCI IDS

CM 2

CRN 195376-89-7
CMF C13 H14 O7

Absolute stereochemistry.



CM 3

CRN 107-21-1
CMF C2 H6 O2


HO-CH₂-CH₂-OH

IT 195376-93-3P 195376-94-4P 195442-95-6P
195540-15-9P

(functionalized bicyclic (meth)acrylates for dental adhesives)

L10 ANSWER 8 OF 14 HCA COPYRIGHT 2003 ACS on STN
127:248519 Polymerizable hybrid monomers. Moszner, Norbert;
Rheinberger, Volker; Zeuner, Frank (Ivoclar Ag, Liechtenstein).
Ger. Offen. DE 19608313 A1 19970828, 13 pp. (German). CODEN:
GWXXBX. APPLICATION: DE 1996-19608313 19960222.

GI



AB The monomers (I; A-E = C:C, C-C; G = H, Me; L = org. connecting group; M = O, NH, CO₂, CONH, O₂CNH; Q = polymerizable group; X = CH₂, O; Y = H, org. group, halogen, NO₂, NH₂, SH; Z = O, NH; M = O, 1; n = 1-4) are obtained by a Diels-Alder reaction of a diene (meth)acrylate with a dienophile followed by attachment of the Q group. I may be used in dentin adhesives. Thus, maleic anhydride was subjected to a Diels-Alder reaction with furfuryl methacrylate and the resulting oxanorbornenedicarboxylic anhydride was monoesterified with allyl alc. to give a monomer which was copolymd. with 2-hydroxyethyl methacrylate to provide a dentin adhesive primer.

IT 195378-02-0P

(dentin adhesive primer; prepn. of hybrid monomers for dentin adhesives)


RN 195378-02-0 HCA

CN 7-Oxabicyclo[2.2.1]hept-5-ene-2,3-dicarboxylic acid, 1-[[[(2-methyl-1-oxo-2-propenyl)oxy]methyl]-, mono-2-propenyl ester, (exo,exo)-, polymer with 2-hydroxyethyl 2-methyl-2-propenoate (9CI) (CA INDEX NAME)

CM 1

CRN 868-77-9

CMF C6 H10 O3



CM 2

CRN 195378-01-9

CMF C16 H18 O7


CCI IDS

CM 3

CRN 194920-51-9

CMF C13 H14 O7


Relative stereochemistry.



CM 4

CRN 107-18-6

CMF C3 H6 O



IT 195540-06-8P 195540-07-9P 195540-08-0P

195540-09-1P 195540-10-4P 195540-11-5P

(prepn. of hybrid monomers for dentin adhesives)

RN 195540-06-8 HCA

CN 7-Oxabicyclo[2.2.1]hept-5-ene-2,3-dicarboxylic acid,
 1-[[[(2-methyl-1-oxo-2-propenyl)oxy]methyl]-, mono[2-
 (ethenyloxy)ethyl] ester, (exo,exo)-, homopolymer (9CI) (CA INDEX
 NAME)

CM 1

CRN 195377-97-0

CMF C17 H20 O8


CCI IDS

CM 2

CRN 194920-51-9

CMF C13 H14 O7


Relative stereochemistry.



CM 3

CRN 764-48-7

CMF C4 H8 O2



RN 195540-07-9 HCA

CN 7-Oxabicyclo[2.2.1]hept-5-ene-2,3-dicarboxylic acid,
1-[[[(2-methyl-1-oxo-2-propenyl)oxy]methyl]-, mono[2-[(1-oxo-2-
propenyl)oxy]methyl] ester, (exo,exo)-, homopolymer (9CI) (CA INDEX
NAME)

CM 1

CRN 195377-98-1

CMF C18 H20 O9


CCI IDS

CM 2

CRN 194920-51-9

CMF C13 H14 O7


Relative stereochemistry.



CM 3

CRN 818-61-1

CMF C5 H8 O3



RN 195540-08-0 HCA

CN 7-Oxabicyclo[2.2.1]hept-5-ene-2,3-dicarboxylic acid,
 1-[[[(2-methyl-1-oxo-2-propenyl)oxy]methyl]-, mono[2-[(2-methyl-1-oxo-2-propenyl)oxy]ethyl] ester, (exo,exo)-, homopolymer (9CI) (CA
 INDEX NAME)

CM 1

CRN 195378-00-8

CMF C19 H22 O9


CCI IDS

CM 2

CRN 194920-51-9

CMF C13 H14 O7


Relative stereochemistry.



CM 3

CRN 868-77-9

CMF C6 H10 O3



RN 195540-09-1 HCA

CN 7-Oxabicyclo[2.2.1]hept-5-ene-2,3-dicarboxylic acid,
1-[[[(2-methyl-1-oxo-2-propenyl)oxy]methyl]-, mono-2-propenyl ester,
(exo,exo)-, homopolymer (9CI) (CA INDEX NAME)

CM 1

CRN 195378-01-9

CMF C16 H18 O7


CCI IDS

CM 2

CRN 194920-51-9

CMF C13 H14 O7


Relative stereochemistry.



CM 3

CRN 107-18-6

CMF C3 H6 O



RN 195540-10-4 HCA

CN 7-Oxabicyclo[2.2.1]hept-5-ene-2,3-dicarboxylic acid,
 1-[[[(2-methyl-1-oxo-2-propenyl)oxy]methyl]-, mono[2-[(2-methyl-1-oxo-
 2-propenyl)oxy]-1-[[[(2-methyl-1-oxo-2-propenyl)oxy]methyl]ethyl]
 ester, (exo,exo)-, homopolymer (9CI) (CA INDEX NAME)

CM 1

CRN 195378-03-1

CMF C24 H28 O11


CCI IDS

CM 2

CRN 194920-51-9


CMF C13 H14 O7

Relative stereochemistry.



CM 3

CRN 1830-78-0
CMF C11 H16 O5



RN 195540-11-5 HCA
CN 7-Oxabicyclo[2.2.1]hept-5-ene-2,3-dicarboxylic acid,
1-[[[(2-methyl-1-oxo-2-propenyl)oxy]methyl]-, monopropyl ester,
(exo,exo)-, homopolymer (9CI) (CA INDEX NAME)


CM 1

CRN 195378-04-2
CMF C16 H20 O7
CCI IDS

CM 2

CRN 194920-51-9
CMF C13 H14 O7


Relative stereochemistry.



CM 3

CRN 71-23-8

CMF C3 H8 O



IT 195378-02-0P

(dentin adhesive primer; prepn. of hybrid monomers for dentin adhesives)

IT 195540-06-8P 195540-07-9P 195540-08-0P

195540-09-1P 195540-10-4P 195540-11-5P

(prepn. of hybrid monomers for dentin adhesives)

L10 ANSWER 9 OF 14 HCA COPYRIGHT 2003 ACS on STN


119:118074 Bicycloheptenedicarboxylic anhydride derivative polymers.

Besecke, Siegmund; Guentherberg, Norbert (BASF A.-G., Germany).

Ger. Offen. DE 4117369 A1 19921203, 7 pp. (German). CODEN: GWXXBX.

APPLICATION: DE 1991-4117369 19910528.

GI



AB Polymers with good resistance to thermal distortion and good flow and color stability are prepd. from the anhydrides I (R = alkyl, alkoxy, aryl, aryloxy, alkylaryl, CN, Cl, Br, CO₂H, carboalkoxy,

acyloxy, carbamoyl; Z = O, S; n = 0-4) and, optionally, .ltoreq.95% comonomer. Peroxide-initiated polymn. of a 23:77 mixt. of I (Z = O, n = 0) and Me methacrylate in THF at 60.degree. for 9 h gave a copolymer with glass temp. 117.degree., decompn. temp. 303.degree., and temp. of 2% wt. loss 263.degree.; vs. 107, 250, and 234, resp., for PMMA.

IT 148976-99-2P, MMA-7-oxabicyclo[4.1.0]hept-5-ene-2,3-dicarboxylic anhydride copolymer 148977-01-9P, MMA-7-oxabicyclo[4.1.0]hept-5-ene-2,3-dicarboxylic anhydride-styrene copolymer

(prepn. of, with good thermal stability)


RN 148976-99-2 HCA

CN 2-Propenoic acid, 2-methyl-, methyl ester, polymer with 3a,4,7,7a-tetrahydro-4,7-epoxyisobenzofuran-1,3-dione (9CI) (CA INDEX NAME)

CM 1

CRN 5426-09-5


CMF C8 H6 O4



CM 2

CRN 80-62-6

CMF C5 H8 O2




RN 148977-01-9 HCA

CN 2-Propenoic acid, 2-methyl-, methyl ester, polymer with ethenylbenzene and 3a,4,7,7a-tetrahydro-4,7-epoxyisobenzofuran-1,3-dione (9CI) (CA INDEX NAME)

CM 1

CRN 5426-09-5


CMF C8 H6 O4



CM 2

CRN 100-42-5


CMF C8 H8



CM 3

CRN 80-62-6

CMF C5 H8 O2



IT 148976-99-2P, MMA-7-oxabicyclo[4.1.0]hept-5-ene-2,3-dicarboxylic anhydride copolymer 148977-01-9P, MMA-7-oxabicyclo[4.1.0]hept-5-ene-2,3-dicarboxylic anhydride-styrene copolymer
(prepn. of, with good thermal stability)

L10 ANSWER 10 OF 14 HCA COPYRIGHT 2003 ACS on STN

118:192255 Immobilized thermolysin and synthesis of precursor of aspartame. Zhou, Qingzhong; Huang, Zhen (Dep. Chem., Peking Univ., Beijing, 100871, Peop. Rep. China). Indian Journal of Chemistry, Section B: Organic Chemistry Including Medicinal Chemistry, 32B(1), 35-9 (English) 1993. CODEN: IJSBDB. ISSN: 0376-4699.

AB Optimal conditions for the thermolysin-catalyzed condensation of Z-DL-Asp-OMe (Z = PhCH2O2C) with H-DL-Phe-OMe to give Z-L-Asp-L-Phe-OMe, the precursor of the synthetic sweetener aspartame were investigated. The immobilized thermolysins were prepd. using 7 polymethacrylate derivs. as carriers by covalent coupling of the polymers with enzyme. The abilities of the immobilized thermolysis to catalyze the condensation reaction were tested and 100% yield was obtained using one of them.

IT 146899-29-8P

(prepn. of, as carrier for thermolysin in peptide coupling)

reactions)


RN 146899-29-8 HCA

CN 2-Propenoic acid, 2-methyl-, 4-[[[(1,3,3a,4,7,7a-hexahydro-1,4-dioxo-4,7-epoxy-2H-isoindol-2-yl)oxy]carbonyl]phenyl ester,
(3a.alpha.,4.alpha.,7.alpha.,7a.alpha.)-, homopolymer (9CI) (CA
INDEX NAME)

CM 1

CRN 146899-28-7

CMF C19 H15 N O7



IT 146899-29-8P

(prepn. of, as carrier for thermolysin in peptide coupling
reactions)

L10 ANSWER 11 OF 14 HCA COPYRIGHT 2003 ACS on STN

114:43679 Oligomerization and copolymerization of endo-N-cyclohexyl-9-oxabicyclo[2.2.1]hept-2-ene-5,6-dicarboximide. Oishi, Tsutomu; Momoi, Masaaki; Fujimoto, Minoru (Tech. Coll., Yamaguchi Univ., Ube, 755, Japan). Polymer Journal (Tokyo, Japan), 22(11), 1007-14 (English) 1990. CODEN: POLJB8. ISSN: 0032-3896.

AB endo-N-Cyclohexyl-9-oxabicyclo[2.2.1]hept-2-ene-5,6-dicarboximide (I) was prepd. from N-cyclohexylmaleimide and furan by Diels-Alder reaction and polymd. The no.-av. mol. wts. (.hivin.Mn) of the polymers insol. in MeOH were 1100-2000, i.e., the d.p. was 4-8. The oligomers having .hivin.Mn <1000 were sol. in MeOH. I was copolymd. with styrene, Me methacrylate, and vinyl acetate in the presence of radical initiators. The monomer reactivity ratios and Alfrey-Price Q-e values were detd.

IT 127122-32-1

(mol. wt. distribution of)


RN 127122-32-1 HCA

CN 2-Propenoic acid, 2-methyl-, methyl ester, polymer with
2-cyclohexyl-3a,4,7,7a-tetrahydro-4,7-epoxy-1H-isoindole-1,3(2H)-
dione (9CI) (CA INDEX NAME)

CM 1

CRN 127122-27-4


CMF C14 H17 N O3



CM 2

CRN 80-62-6


CMF C5 H8 O2



IT 127122-32-1
(mol. wt. distribution of)

L10 ANSWER 12 OF 14 HCA COPYRIGHT 2003 ACS on STN
112:236040 Heat-resistant oxabicycloheptenedicarboximide copolymers.
Kinoshita, Seigo; Kato, Kenji; Haruta, Yukinori; Oishi, Tsutomu
(Nippon Oils and Fats Co., Ltd., Japan). Jpn. Kokai Tokkyo Koho JP
01311112 A2 19891215 Heisei, 4 pp. (Japanese). CODEN: JKXXAF.
APPLICATION: JP 1988-140540 19880609.

GI



AB The title copolymers are composed of oxabicycloheptenedicarboximides I (R = H, C1-18 alkyl, cycloalkyl, aryl, aralkyl) and other vinyl monomers. Thus, reflux of 17.6 g N-cyclohexylmaleimide and 20.4 g furan in toluene for 3 h gave 75.0% I (R = cyclohexyl), 0.372 g of which was polymd. with 0.628 g styrene in presence of Perhexyne 25B at 120.degree. for 20 h to give 88.7% copolymer with no.-av. mol. wt. 10.51 .times. 104, wt.-av. mol. wt. 12.19 .times. 104, and softening point 148.degree..

IT 127122-32-1P

(prepn. of, heat-resistant)


RN 127122-32-1 HCA

CN 2-Propenoic acid, 2-methyl-, methyl ester, polymer with
2-cyclohexyl-3a,4,7,7a-tetrahydro-4,7-epoxy-1H-isoindole-1,3(2H)-
dione (9CI) (CA INDEX NAME)

CM 1

CRN 127122-27-4


CMF C14 H17 N O3



CM 2

CRN 80-62-6

CMF C5 H8 O2



IT 127122-32-1P

(prepn. of, heat-resistant)

L10 ANSWER 13 OF 14 HCA COPYRIGHT 2003 ACS on STN

81:78318 Radical polymerization of furan with maleic anhydride through
the Diels-Alder adduct. Kamo, Bunzo; Morita, Isamu; Horie, Satoshi;
Furusawa, Shijo (Fac. Sci. Eng., Chuo Univ., Tokyo, Japan). Polymer
Journal (Tokyo, Japan), 6(2), 121-31 (English) 1974. CODEN: POLJB8.
ISSN: 0032-3896.AB An equimolar reaction of furan, 2-methylfuran, or 2,5-dimethylfuran
with maleic anhydride (I) in acetone at 60.deg. gave the Diels-Alder
exo adducts (II, R1, R2 = H, Me) which were polymd. at 70.deg. in
the presence of azobisisobutyronitrile as radical initiator. The
reaction of furfuryl methacrylate with I gave the same result.

IT 52520-31-7P

(prepn. of)

RN 52520-31-7 HCA

CN 2-Propenoic acid, 2-methyl-, (3,3a,7,7a-tetrahydro-1,3-dioxo-4,7-
epoxyisobenzofuran-4(1H)-yl)methyl ester,
(3a.alpha.,4.beta.,7.beta.,7a.alpha.)-, homopolymer (9CI) (CA INDEX


NAME)

CM 1

CRN 52485-74-2

CMF C13 H12 O6

Relative stereochemistry.



IT 52520-31-7P
(prepn. of)

L10 ANSWER 14 OF 14 HCA COPYRIGHT 2003 ACS on STN

66:46639 Polymers of adducts of furfuryl methacrylate with maleic anhydride and maleimide. Mikhailov, Marin; Budevskia, Khr.; Berlin, A. A. Comptes Rendus de l'Academie Bulgare des Sciences, 19(11), 1019-22 (English) 1966. CODEN: CRABAA.

AB A furfuryl methacrylate-maleic anhydride adduct and a furfuryl methacrylate-maleimide adduct were homopolymd. in 10% C6H6 and AcOMe solns. in the presence of 2% azobisisobutyronitrile to yield yellowish to white polymers which swell in C6H6, AcOMe, CHCl3, and HCONMe2 and do not melt up to 280.degree.. The polymers, which are assumed to be partially crosslinked polyadducts, have nearly identical residues when heated above 293.degree.. Under 100.degree., they undergo deadduction, and the .pi.-complexes formed dissociate from 100 to 293.degree., while part of the remaining dienophile is polymd. and part is volatilized.

IT 31343-32-5P 31343-33-6P
(prepn. of)


RN 31343-32-5 HCA

CN 2-Propenoic acid, 2-methyl-, (3,3a,7,7a-tetrahydro-1,3-dioxo-4,7-epoxyisobenzofuran-4(1H)-yl)methyl ester, homopolymer (9CI) (CA INDEX NAME)

CM 1

CRN 7726-38-7

CMF C13 H12 O6




RN 31343-33-6 HCA
 CN Methacrylic acid, ester with 1-(hydroxymethyl)-7-oxabicyclo[2.2.1]hept-5-ene-2,3-dicarboximide, polymers (8CI) (CA INDEX NAME)

CM 1

CRN 7726-39-8

CMF C13 H13 N O5



IT 31343-32-5P 31343-33-6P
 (prepn. of)